


Screening Level Risk Assessment Package

Esso Gettysburg



ENVIRONMENTAL
RESEARCH
CONSULTING

National Oceanic and
Atmospheric Administration

Office of National Marine Sanctuaries
Daniel J. Basta, Director
Lisa Symons
John Wagner

Office of Response and Restoration
Dave Westerholm, Director
Debbie Payton
Doug Helton

Photo: Photograph of *Esso Gettysburg*
Source: <http://www.uboot.net/allies/merchants/ships/2949.html>



Table of Contents

Project Background	ii
Executive Summary	1
Section 1: Vessel Background Information: Remediation of Underwater Legacy	
Environmental Threats (RULET).....	2
Vessel Particulars.....	2
Casualty Information.....	3
Wreck Location.....	4
Casualty Narrative	4
General Notes	5
Wreck Condition/Salvage History	5
Archaeological Assessment	5
Assessment.....	5
Background Information References	6
Vessel Risk Factors.....	7
Section 2: Environmental Impact Modeling.....	13
Release Scenarios Used in the Modeling	13
Oil Type for Release	14
Oil Thickness Thresholds.....	14
Potential Impacts to the Water Column.....	15
Potential Water Surface Slick.....	16
Potential Shoreline Impacts.....	19
Section 3: Ecological Resources At Risk	22
Ecological Risk Factors	24
Section 4: Socio-Economic Resources At Risk	29
Socio-Economic Risk Factors	32
Section 5: Overall Risk Assessment and Recommendations for Assessment,	
Monitoring, or Remediation	37

Project Background

The past century of commerce and warfare has left a legacy of thousands of sunken vessels along the U.S. coast. Many of these wrecks pose environmental threats because of the hazardous nature of their cargoes, presence of munitions, or bunker fuel oils left onboard. As these wrecks corrode and decay, they may release oil or hazardous materials. Although a few vessels, such as USS *Arizona* in Hawaii, are well-publicized environmental threats, most wrecks, unless they pose an immediate pollution threat or impede navigation, are left alone and are largely forgotten until they begin to leak.

In order to narrow down the potential sites for inclusion into regional and area contingency plans, in 2010, Congress appropriated \$1 million to identify the most ecologically and economically significant potentially polluting wrecks in U.S. waters. This project supports the U.S. Coast Guard and the Regional Response Teams as well as NOAA in prioritizing threats to coastal resources while at the same time assessing the historical and cultural significance of these nonrenewable cultural resources.

The potential polluting shipwrecks were identified through searching a broad variety of historical sources. NOAA then worked with Research Planning, Inc., RPS ASA, and Environmental Research Consulting to conduct the modeling forecasts, and the ecological and environmental resources at risk assessments.

Initial evaluations of shipwrecks located within American waters found that approximately 600-1,000 wrecks could pose a substantial pollution threat based on their age, type and size. This includes vessels sunk after 1891 (when vessels began being converted to use oil as fuel), vessels built of steel or other durable material (wooden vessels have likely deteriorated), cargo vessels over 1,000 gross tons (smaller vessels would have limited cargo or bunker capacity), and any tank vessel.

Additional ongoing research has revealed that 87 wrecks pose a potential pollution threat due to the violent nature in which some ships sank and the structural reduction and demolition of those that were navigational hazards. To further screen and prioritize these vessels, risk factors and scores have been applied to elements such as the amount of oil that could be on board and the potential ecological or environmental impact.

Executive Summary: *Esso Gettysburg*

The tanker *Esso Gettysburg*, torpedoed and sunk during World War II off the coast of Georgia in 1943, was identified as a potential pollution threat, thus a screening-level risk assessment was conducted. The different sections of this document summarize what is known about the *Esso Gettysburg*, the results of environmental impact modeling composed of different release scenarios, the ecological and socio-economic resources that would be at risk in the event of releases, the screening-level risk scoring results and overall risk assessment, and recommendations for assessment, monitoring, or remediation.



Based on this screening-level assessment, each vessel was assigned a summary score calculated using the seven risk criteria described in this report. For the Worst Case Discharge, *Esso Gettysburg* scores High with 18 points; for the Most Probable Discharge (10% of the Worst Case volume), *Esso Gettysburg* also scores High with 16 points. Given the high scores for both Worst Case Discharge and a Most Probable Discharge for *Esso Gettysburg*, NOAA would typically recommend that this site be considered for further assessment to determine the vessel condition, amount of oil onboard, and feasibility of oil removal action. However, the archival and archaeological review suggests that the vessel may have incurred substantial structural damage. Given the nature of the casualty and uncertainty of the vessel condition and location, NOAA recommends that surveys of opportunity with state, federal, or academic entities be used to attempt to locate this vessel. Also, general notations should be made in the Area Contingency Plans so that if a mystery spill is reported in the general area, this vessel could be investigated as a source. Outreach efforts with commercial and recreational fishermen who frequent the area would be helpful to gain awareness of localized spills in the general area where the vessel is believed lost.

Vessel Risk Factors		Risk Score	
Pollution Potential Factors	A1: Oil Volume (total bbl)	Med	
	A2: Oil Type		
	B: Wreck Clearance		
	C1: Burning of the Ship		
	C2: Oil on Water		
	D1: Nature of Casualty		
Archaeological Assessment	D2: Structural Breakup	Not Scored	
	Archaeological Assessment		
Operational Factors	Wreck Orientation	Not Scored	
	Depth		
	Visual or Remote Sensing Confirmation of Site Condition		
	Other Hazardous Materials Onboard		
	Munitions Onboard		
	Gravesite (Civilian/Military)		
		WCD	MP (10%)
Ecological Resources	3A: Water Column Resources	Med	Med
	3B: Water Surface Resources	High	High
	3C: Shore Resources	Med	Low
Socio-Economic Resources	4A: Water Column Resources	High	Med
	4B: Water Surface Resources	High	High
	4C: Shore Resources	High	High
Summary Risk Scores		18	16

The determination of each risk factor is explained in the document. This summary table is found on page 38.

SECTION 1: VESSEL BACKGROUND INFORMATION: REMEDIATION OF UNDERWATER LEGACY ENVIRONMENTAL THREATS (RULET)

Vessel Particulars

Official Name: *Esso Gettysburg*

Official Number: 241409

Vessel Type: Tanker

Vessel Class: T2-SE-A1 Type Tanker

Former Names: *Gettysburg*

Year Built: 1942

Builder: Sun Shipbuilding & Dry Dock Company (Sun Ship), Chester, PA

Builder's Hull Number: 241

Flag: American

Owner at Loss: Standard Oil Co. of New Jersey

Controlled by: Unknown

Chartered to: United States War Shipping Administration (Not on Govt. mission when lost)

Operated by: Standard Oil Co. of New Jersey

Homeport: Wilmington, DE

Length: 503 feet

Beam: 68 feet

Depth: 39 feet

Gross Tonnage: 10,173

Net Tonnage: 6,134

Hull Material: Steel

Hull Fastenings: Welded

Powered by: Oil-fired steam

Bunker Type: Heavy fuel oil (Bunker C)


Bunker Capacity (bbl): 11,026

Average Bunker Consumption (bbl) per 24 hours: 295

Liquid Cargo Capacity (bbl): 138,330

Dry Cargo Capacity: 15,200 cubic feet

Tank or Hold Description: Vessel had 9 cargo tanks, tanks Nos. 2 to 9 being divided by two longitudinal bulkheads so that there were 2 wing tanks and a center tank; tank No. 1 was divided by a center line bulkhead into 2 tanks. 1 forward deep tank split starboard & port



Casualty Information

Port Departed: Port Arthur, TX

Destination Port: Philadelphia, PA

Date Departed: June 6, 1943

Date Lost: June 10, 1943

Number of Days Sailing: ≈ 5

Cause of Sinking: Act of War (Torpedoes)

Latitude (DD): 31.00023

Longitude (DD): -79.24976

Nautical Miles to Shore: 106

Nautical Miles to NMS: 97

Nautical Miles to MPA: 0

Nautical Miles to Fisheries: Unknown

Approximate Water Depth (Ft): 2,300

Bottom Type: Sand

Is There a Wreck at This Location? Unknown, the wreck has never been located

Wreck Orientation: Unknown

Vessel Armament: Vessel was armed with one 4in gun, one 3in gun, and eight 20mm guns

Cargo Carried when Lost: 120,120 bbl of West Texas crude oil (API gravity 35) loaded in tanks 2 to 7 inclusive, tank 8 across, and 9 center

Cargo Oil Carried (bbl): 120,120

Cargo Oil Type: West Texas Crude (API gravity 35)

Probable Fuel Oil Remaining (bbl): $\leq 11,026$

Fuel Type: Heavy fuel oil (Bunker C)

Total Oil Carried (bbl): $\leq 131,146$

Dangerous Cargo or Munitions: Yes

Munitions Carried: Munitions for onboard weapons

Demolished after Sinking: No

Salvaged: No

Cargo Lost: Yes

Reportedly Leaking: No

Historically Significant: Yes

Gravesite: Yes

Salvage Owner: Not known if any

Wreck Location




Chart Number: 11009

Casualty Narrative

"At 20.00 hours on 10 Jun, 1943, the unescorted *Esso Gettysburg* (Master Peder A. Johnson, lost) was hit by two torpedoes from *U-66* about 100 miles southeast of Savannah, Georgia, shortly after she received a U-boat warning, steaming on a zigzag course at 15.5 knots. One torpedo struck the port side between the #6 and #7 tanks, ripped up 25 feet of deck, blew oil 100 feet into the air and disabled the steering gear. Seconds later the second struck on the port side at the engine room, causing an immediate fire as she began to settle by the stern and listed to port. Oil from the two tanks was spread into the water and was ignited by the second explosion. The flames spread 100 feet on both sides, while smoke rose over 1000 feet in the air. The eight officers, 37 men and 27 armed guards (the ship was armed with one 4in, one 3in and eight 20mm guns) attempted to launch some lifeboats, but failed because of the intense flames. Only 15 men (seven armed guards, three officers and five crewmen) survived because they jumped overboard and swam away as fast as they could. The entirely submerged tanker, except for a small part of the bow, was last seen about 03.00 hours on 12 June and eventually sank. The survivors found a badly burned lifeboat after swimming for three hours and extinguished the fire. All were picked up by the American steam passenger ship *George Washington* the next day after they were sighted by an U.S. Army patrol aircraft and landed in Charleston, South Carolina, the same day. The armed guards ensign was awarded the Navy Cross."

<http://www.uboa.net/allies/merchants/ships/2949.html>

General Notes

NOAA Automated Wreck and Obstruction Information System (AWOIS) Data:

DESCRIPTION-

NO.1162; TANKER, 10172 GT; SUNK 6/10/42 BY SUBMARINE; POSITION ACCURACY 3-5 MILES. 12/30/50

SURVEY REQUIREMENT NOT DETERMINED.

Vessel was not on Government Service but was carrying an Atlantic Refining Co., cargo.

Wreck Condition/Salvage History

Unknown; the wreck has never been located or surveyed.

Archaeological Assessment

The archaeological assessment provides additional primary source based documentation about the sinking of vessels. It also provides condition-based archaeological assessment of the wrecks when possible. It does not provide a risk-based score or definitively assess the pollution risk or lack thereof from these vessels, but includes additional information that could not be condensed into database form.

Where the current condition of a shipwreck is not known, data from other archaeological studies of similar types of shipwrecks provide the means for brief explanations of what the shipwreck might look like and specifically, whether it is thought there is sufficient structural integrity to retain oil. This is more subjective than the Pollution Potential Tree and computer-generated resource at risk models, and as such provides an additional viewpoint to examine risk assessments and assess the threat posed by these shipwrecks. It also addresses questions of historical significance and the relevant historic preservation laws and regulations that will govern on-site assessments.

In some cases where little additional historic information has been uncovered about the loss of a vessel, archaeological assessments cannot be made with any degree of certainty and were not prepared. For vessels with full archaeological assessments, NOAA archaeologists and contracted archivists have taken photographs of primary source documents from the National Archives that can be made available for future research or on-site activities.

Assessment

The wreck of *Esso Gettysburg* has never been located so there are no site reports that would allow NOAA archaeologists to provide a condition based archaeological assessment of the shipwreck. Some additional analysis can be made based on the historic sinking reports of the ship that may be of utility to the U.S. Coast Guard. We know from archival research that two torpedoes struck the ship with devastating results.

The official Summary of Statements by Survivors states that the first torpedo “struck port side between midships house and mainmasts at #6 or #7 tank (Fig. 1-1), exploding on contact, rupturing 25’ of deck and hull near sheer strake and raising 100’ geyser of oil and water spray, tearing up deck over #6 and #7 port wing tanks, knocking longitudinal bulkhead down, disabling steering apparatus. A second torpedo entered port side 3-5 seconds later at engine room, exploding on contact, causing an immediate fire that

enveloped after part of ship in flames. Within a few seconds the fire spread forward to midships house. Flame spread on water 50' to 100' on both sides of ship."




Figure 1-1: U.S. Coast Guard diagram of the location of torpedo impacts on *Esso Gettysburg* (Image courtesy of National Archives, Washington, DC)

Based on the location of the torpedo impacts, it is likely that the ship's bunker tanks by the engine room were destroyed and that all aft tanks were damaged or breached by the flames before the vessel sank. Although the condition of the forward tanks is not known, ongoing research strongly suggests that vessels in great depths of water are generally found in an upright orientation. This orientation has often lead to loss of oil from vents and piping long before loss of structural integrity of hull plates from corrosion or other physical impacts. As it is believed that this vessel is in water greater than 2,000 feet, it is likely to have settled upright and may no longer contain oil.

The only way to conclusively determine the condition of the shipwreck, however, will be to examine the site after it is discovered. Should the vessel be located in a survey of opportunity or due to a mystery spill attributed to this vessel, it should be noted that this vessel is of historic significance and will require appropriate actions be taken under the National Historic Preservation Act (NHPA) and the Sunken Military Craft Act (SMCA) prior to any actions that could impact the integrity of the vessel. This vessel may be eligible for listing on the National Register of Historic Places. The site is also considered a war grave and appropriate actions should be undertaken to minimize disturbance to the site.

Background Information References

Vessel Image Sources: <http://www.uboaat.net/allies/merchants/ships/2949.html>

Construction Diagrams or Plans in RULET Database? No, but ONMS has capacity plans for a T2-SE-A1 Type Tanker

Text References:

-AWOIS database

-<http://www.uboaat.net/allies/merchants/ships/2949.html>

Vessel Risk Factors

In this section, the risk factors that are associated with the vessel are defined and then applied to the *Esso Gettysburg* based on the information available. These factors are reflected in the pollution potential risk assessment development by the U.S. Coast Guard Salvage Engineering Response Team (SERT) as a means to apply a salvage engineer's perspective to the historical information gathered by NOAA. This analysis reflected in Figure 1-2 is simple and straightforward and, in combination with the accompanying archaeological assessment, provides a picture of the wreck that is as complete as possible based on current knowledge and best professional judgment. This assessment does not take into consideration operational constraints such as depth or unknown location, but rather attempts to provide a replicable and objective screening of the historical data for each vessel. SERT reviewed the general historical information available for the database as a whole and provided a stepwise analysis for an initial indication of Low/Medium/High values for each vessel.




Figure 1-2: U.S. Coast Guard Salvage Engineering Response Team (SERT) developed the above Pollution Potential Decision Tree.

In some instances, nuances from the archaeological assessment may provide additional input that will amend the score for Section 1. Where available, additional information that may have bearing on operational considerations for any assessment or remediation activities is provided.

Each risk factor is characterized as High, Medium, or Low Risk or a category-appropriate equivalent such as No, Unknown, Yes, or Yes Partially. The risk categories correlate to the decision points reflected in Figure 1-2.

Each of the risk factors also has a “data quality modifier” that reflects the completeness and reliability of the information on which the risk ranks were assigned. The quality of the information is evaluated with respect to the factors required for a reasonable preliminary risk assessment. The data quality modifier scale is:

- **High Data Quality:** All or most pertinent information on wreck available to allow for thorough risk assessment and evaluation. The data quality is high and confirmed.
- **Medium Data Quality:** Much information on wreck available, but some key factor data are missing or the data quality is questionable or not verified. Some additional research needed.
- **Low Data Quality:** Significant issues exist with missing data on wreck that precludes making preliminary risk assessment, and/or the data quality is suspect. Significant additional research needed.

In the following sections, the definition of low, medium, and high for each risk factor is provided. Also, the classification for the *Esso Gettysburg* is provided, both as text and as shading of the applicable degree of risk bullet.

Pollution Potential Factors

Risk Factor A1: Total Oil Volume

The oil volume classifications correspond to the U.S. Coast Guard spill classifications:

- **Low Volume: Minor Spill** <240 bbl (10,000 gallons)
- **Medium Volume: Medium Spill** ≥240 – 2,400 bbl (100,000 gallons)
- **High Volume: Major Spill** ≥2,400 bbl (≥100,000 gallons)

The oil volume risk classifications refer to the volume of the most-likely Worst Case Discharge from the vessel and are based on the amount of oil believed or confirmed to be on the vessel.

The *Esso Gettysburg* is ranked as High Volume because it is thought to have a potential for up to 131,146 bbl, although some of that was lost at the time of the casualty due to the explosion and fire. This volume estimate takes into account the amount of time the vessel was underway prior to the casualty. Data quality is medium.

The risk factor for volume also incorporates any reports or anecdotal evidence of actual leakage from the vessel or reports from divers of oil in the overheads, as opposed to potential leakage. This reflects the history of the vessel’s leakage. There are no reports of leakage from the *Esso Gettysburg*.

Risk Factor A2: Oil Type

The oil type(s) on board the wreck are classified only with regard to persistence, using the U.S. Coast Guard oil grouping¹. (Toxicity is dealt with in the impact risk for the Resources at Risk classifications.)

The three oil classifications are:

- **Low Risk: Group I Oils** – non-persistent oil (e.g., gasoline)
- **Medium Risk: Group II – III Oils** – medium persistent oil (e.g., diesel, No. 2 fuel, light crude, medium crude)
- **High Risk: Group IV** – high persistent oil (e.g., heavy crude oil, No. 6 fuel oil, Bunker C)

The *Esso Gettysburg* is classified as Medium Risk because the cargo is medium crude oil, a Group III oil type. Data quality is high.

Was the wreck demolished?

Risk Factor B: Wreck Clearance

This risk factor addresses whether or not the vessel was historically reported to have been demolished as a hazard to navigation or by other means such as depth charges or aerial bombs. This risk factor is based on historic records and does not take into account what a wreck site currently looks like. The risk categories are defined as:

- **Low Risk:** The wreck was reported to have been entirely destroyed after the casualty
- **Medium Risk:** The wreck was reported to have been partially cleared or demolished after the casualty
- **High Risk:** The wreck was not reported to have been cleared or demolished after the casualty
- **Unknown:** It is not known whether or not the wreck was cleared or demolished at the time of or after the casualty

The *Esso Gettysburg* is classified as High Risk because there are no known historic accounts of the wreck being demolished as a hazard to navigation. Data quality is high.

Was significant cargo or bunker lost during casualty?

Risk Factor C1: Burning of the Ship

This risk factor addresses any burning that is known to have occurred at the time of the vessel casualty and may have resulted in oil products being consumed or breaks in the hull or tanks that would have increased the potential for oil to escape from the shipwreck. The risk categories are:

- **Low Risk:** Burned for multiple days
- **Medium Risk:** Burned for several hours
- **High Risk:** No burning reported at the time of the vessel casualty
- **Unknown:** It is not known whether or not the vessel burned at the time of the casualty

¹ Group I Oil or Nonpersistent oil is defined as “a petroleum-based oil that, at the time of shipment, consists of hydrocarbon fractions: At least 50% of which, by volume, distill at a temperature of 340°C (645°F); and at least 95% of which, by volume, distill at a temperature of 370°C (700°F).”

Group II - Specific gravity less than 0.85 crude [API° >35.0]

Group III - Specific gravity between 0.85 and less than .95 [API° ≤35.0 and >17.5]

Group IV - Specific gravity between 0.95 to and including 1.0 [API° ≤17.5 and >10.0]

The *Esso Gettysburg* is classified as Medium Risk because the ship burned prior to sinking. Data quality is high.

Risk Factor C2: Reported Oil on the Water

This risk factor addresses reports of oil on the water at the time of the vessel casualty. The amount is relative and based on the number of available reports of the casualty. Seldom are the reports from trained observers so this is very subjective information. The risk categories are defined as:

- **Low Risk:** Large amounts of oil reported on the water by multiple sources
- **Medium Risk:** Moderate to little oil reported on the water during or after the sinking event
- **High Risk:** No oil reported on the water
- **Unknown:** It is not known whether or not there was oil on the water at the time of the casualty

The *Esso Gettysburg* is classified as Low Risk because the oil was reported to have spread across the water as the vessel went down. Data quality is high.

Is the cargo area damaged?

Risk Factor D1: Nature of the Casualty

This risk factor addresses the means by which the vessel sank. The risk associated with each type of casualty is determined by the how violent the sinking event was and the factors that would contribute to increased initial damage or destruction of the vessel (which would lower the risk of oil, other cargo, or munitions remaining on board). The risk categories are:

- **Low Risk:** Multiple torpedo detonations, multiple mines, severe explosion
- **Medium Risk:** Single torpedo, shellfire, single mine, rupture of hull, breaking in half, grounding on rocky shoreline
- **High Risk:** Foul weather, grounding on soft bottom, collision
- **Unknown:** The cause of the loss of the vessel is not known

The *Esso Gettysburg* is classified as Low Risk because there were two torpedo detonations, explosions, and fire. Data quality is high.

Risk Factor D2: Structural Breakup

This risk factor takes into account how many pieces the vessel broke into during the sinking event or since sinking. This factor addresses how likely it is that multiple components of a ship were broken apart including tanks, valves, and pipes. Experience has shown that even vessels broken in three large sections can still have significant pollutants on board if the sections still have some structural integrity. The risk categories are:

- **Low Risk:** The vessel is broken into more than three pieces
- **Medium Risk:** The vessel is broken into two-three pieces
- **High Risk:** The vessel is not broken and remains as one contiguous piece
- **Unknown:** It is currently not known whether or not the vessel broke apart at the time of loss or after sinking

The *Esso Gettysburg* is classified as Unknown because this vessel was in one piece at the time of the casualty, and has not been located since; whether additional structural breakup occurred is unknown as location is unknown. Data quality is high.

Factors That May Impact Potential Operations

Orientation (degrees)

This factor addresses what may be known about the current orientation of the intact pieces of the wreck (with emphasis on those pieces where tanks are located) on the seafloor. For example, if the vessel turtled, not only may it have avoided demolition as a hazard to navigation, but it has a higher likelihood of retaining an oil cargo in the non-vented and more structurally robust bottom of the hull.

The location of the *Esso Gettysburg* is unknown. Data quality is low.

Depth

Depth information is provided where known. In many instances, depth will be an approximation based on charted depths at the last known locations.

The depth for *Esso Gettysburg* is believed to be greater than 2,300 feet due to the last known location. Data quality is low.

Visual or Remote Sensing Confirmation of Site Condition

This factor takes into account what the physical status of wreck site as confirmed by remote sensing or other means such as ROV or diver observations and assesses its capability to retain a liquid cargo. This assesses whether or not the vessel was confirmed as entirely demolished as a hazard to navigation, or severely compromised by other means such as depth charges, aerial bombs, or structural collapse.

The location of the *Esso Gettysburg* is unknown. Data quality is low.

Other Hazardous (Non-Oil) Cargo on Board

This factor addresses hazardous cargo other than oil that may be on board the vessel and could potentially be released, causing impacts to ecological and socio-economic resources at risk.

There are no reports of hazardous materials onboard. Data quality is high.

Munitions on Board

This factor addresses hazardous cargo other than oil that may be on board the vessel and could potentially be released or detonated causing impacts to ecological and socio-economic resources at risk.

The *Esso Gettysburg* had munitions for onboard weapons, one 5-inch gun, one 3-inch gun, and eight 20-mm guns. Data quality is high.

Vessel Pollution Potential Summary

Table 1-1 summarizes the risk factor scores for the pollution potential and mitigating factors that would reduce the pollution potential for the *Esso Gettysburg*. Operational factors are listed but do not have a risk score.

Table 1-1: Summary matrix for the vessel risk factors for the *Esso Gettysburg* color-coded as red (high risk), yellow (medium risk), and green (low risk).

Vessel Risk Factors		Data Quality Score	Comments	Risk Score
Pollution Potential Factors	A1: Oil Volume (total bbl)	Medium	Maximum of 131,146 bbl, no known leakage at this time	Med
	A2: Oil Type	High	Cargo is West Texas crude oil, a Group III oil type	
	B: Wreck Clearance	High	No	
	C1: Burning of the Ship	High	Yes	
	C2: Oil on Water	High	Yes	
	D1: Nature of Casualty	High	Two torpedoes, explosion and fire	
	D2: Structural Breakup	High	Unknown	
Archaeological Assessment	Archaeological Assessment	High	Detailed sinking reports exist, assessment is believed to be very accurate	Not Scored
Operational Factors	Wreck Orientation	Low	Unknown	Not Scored
	Depth	Low	Unknown	
	Visual or Remote Sensing Confirmation of Site Condition	Low	Unknown	
	Other Hazardous Materials Onboard	High	No	
	Munitions Onboard	High	one 4in gun, one 3in gun, eight 20mm guns	
	Gravesite (Civilian/Military)	High	Yes	
	Historical Protection Eligibility (NHPA/SMCA)	High	NHPA and possibly SMCA	

SECTION 2: ENVIRONMENTAL IMPACT MODELING

To help evaluate the potential transport and fates of releases from sunken wrecks, NOAA worked with RPS ASA to run a series of generalized computer model simulations of potential oil releases. The results are used to assess potential impacts to ecological and socio-economic resources, as described in Sections 3 and 4. The modeling results are useful for this screening-level risk assessment; however, it should be noted that detailed site/vessel/and seasonally specific modeling would need to be conducted prior to any intervention on a specific wreck.

Release Scenarios Used in the Modeling

The potential volume of leakage at any point in time will tend to follow a probability distribution. Most discharges are likely to be relatively small, though there could be multiple such discharges. There is a lower probability of larger discharges, though these scenarios would cause the greatest damage. A **Worst Case Discharge** (WCD) would involve the release of all of the cargo oil and bunkers present on the vessel. In the case of the *Esso Gettysburg* this would be 132,000 bbl (rounded up from 131,146 bbl) based on current estimates of the maximum amount of oil remaining onboard the wreck.

The likeliest scenario of oil release from most sunken wrecks, including the *Esso Gettysburg*, is a small, episodic release that may be precipitated by disturbance of the vessel in storms. Each of these episodic releases may cause impacts and require a response. **Episodic** releases are modeled using 1% of the WCD. Another scenario is a very low chronic release, i.e., a relatively regular release of small amounts of oil that causes continuous oiling and impacts over the course of a long period of time. This type of release would likely be precipitated by corrosion of piping that allows oil to flow or bubble out at a slow, steady rate. **Chronic** releases are modeled using 0.1% of the WCD.

The **Most Probable** scenario is premised on the release of all the oil from one tank. In the absence of information on the number and condition of the cargo or fuel tanks for all the wrecks being assessed, this scenario is modeled using 10% of the WCD. The **Large** scenario is loss of 50% of the WCD. The five major types of releases are summarized in Table 2-1. The actual type of release that occurs will depend on the condition of the vessel, time factors, and disturbances to the wreck. Note that, the episodic and chronic release scenarios represent a small release that is repeated many times, potentially repeating the same magnitude and type of impact(s) with each release. The actual impacts would depend on the environmental factors such as real-time and forecast winds and currents during each release and the types/quantities of ecological and socio-economic resources present.

The model results here are based on running the RPS ASA Spill Impact Model Application Package (SIMAP) two hundred times for each of the five spill volumes shown in Table 2-1. The model randomly selects the date of the release, and corresponding environmental, wind, and ocean current information from a long-term wind and current database.

When a spill occurs, the trajectory, fate, and effects of the oil will depend on environmental variables, such as the wind and current directions over the course of the oil release, as well as seasonal effects. The magnitude and nature of potential impacts to resources will also generally have a strong seasonal component (e.g., timing of bird migrations, turtle nesting periods, fishing seasons, and tourism seasons).

Table 2-1: Potential oil release scenario types for the *Esso Gettysburg*.

Scenario Type	Release per Episode	Time Period	Release Rate	Relative Likelihood	Response Tier
Chronic (0.1% of WCD)	132 bbl	Fairly regular intervals or constant	100 bbl over several days	More likely	Tier 1
Episodic (1% of WCD)	1,320 bbl	Irregular intervals	Over several hours or days	Most Probable	Tier 1-2
Most Probable (10% of WCD)	13,200 bbl	One-time release	Over several hours or days	Most Probable	Tier 2
Large (50% of WCD)	66,000 bbl	One-time release	Over several hours or days	Less likely	Tier 2-3
Worst Case	132,000 bbl	One-time release	Over several hours or days	Least likely	Tier 3

The modeling results represent 200 simulations for each spill volume with variations in spill trajectory based on winds and currents. The spectrum of the simulations gives a perspective on the variations in likely impact scenarios. Some resources will be impacted in nearly all cases; some resources may not be impacted unless the spill trajectory happens to go in that direction based on winds and currents at the time of the release and in its aftermath.

For the large and WCD scenarios, the duration of the release was assumed to be 12 hours, envisioning a storm scenario where the wreck is damaged or broken up, and the model simulations were run for a period of 30 days. The releases were assumed to be from a depth between 2-3 meters above the sea floor, using the information known about the wreck location and depth. It is important to acknowledge that these scenarios are only for this screening-level assessment. Detailed site/vessel/and seasonally specific modeling would need to be conducted prior to any intervention on a specific wreck.

Oil Type for Release

The *Esso Gettysburg* contained a maximum of 120,120 bbl of West Texas crude oil (a Group III oil) as cargo and 11,026 bbl of bunker fuel oil (a Group IV oil). Because the bulk of the oil likely remaining on board is West Texas crude oil, the oil spill model was run using medium crude oil.

Oil Thickness Thresholds

The model results are reported for different oil thickness thresholds, based on the amount of oil on the water surface or shoreline and the resources potentially at risk. Table 2-2 shows the terminology and thicknesses used in this report, for both oil thickness on water and the shoreline. For oil on the water surface, a thickness of 0.01 g/m², which would appear as a barely visible sheen, was used as the threshold for socio-economic impacts because often fishing is prohibited in areas with any visible oil, to prevent contamination of fishing gear and catch. A thickness of 10 g/m² was used as the threshold for ecological impacts, primarily due to impacts to birds, because that amount of oil has been observed to be enough to mortally impact birds and other wildlife. In reality, it is very unlikely that oil would be evenly distributed on the water surface. Spilled oil is always distributed patchily on the water surface in bands or tarballs with clean water in between. So, Table 2-2a shows the number of tarballs per acre on the water surface for these oil thickness thresholds, assuming that each tarball was a sphere that was 1 inch in diameter.

For oil stranded onshore, a thickness of 1 g/m² was used as the threshold for socio-economic impacts because that amount of oil would conservatively trigger the need for shoreline cleanup on amenity beaches. A thickness of 100 g/m² was used as the threshold for ecological impacts based on a synthesis of the literature showing that shoreline life has been affected by this degree of oiling.² Because oil often strands onshore as tarballs, Table 2-2b shows the number of tarballs per m² on the shoreline for these oil thickness thresholds, assuming that each tarball was a sphere that was 1 inch in diameter.

Table 2-2a: Oil thickness thresholds used in calculating area of water impacted. Refer to Sections 3 and 4 for explanations of the thresholds for ecological and socio-economic resource impacts.

Oil Description	Sheen Appearance	Approximate Sheen Thickness		No. of 1 inch Tarballs	Threshold/Risk Factor
Oil Sheen	Barely Visible	0.00001 mm	0.01 g/m ²	~5-6 tarballs per acre	Socio-economic Impacts to Water Surface/Risk Factor 4B-1 and 2
Heavy Oil Sheen	Dark Colors	0.01 mm	10 g/m ²	~5,000-6,000 tarballs per acre	Ecological Impacts to Water Surface/ Risk Factor 3B-1 and 2

Table 2-2b: Oil thickness thresholds used in calculating miles of shoreline impacted. Refer to Sections 3 and 4 for explanations of the thresholds for ecological and socio-economic resource impacts.

Oil Description	Oil Appearance	Approximate Sheen Thickness		No. of 1 inch Tarballs	Threshold/Risk Factor
Oil Sheen/Tarballs	Dull Colors	0.001 mm	1 g/m ²	~0.12-0.14 tarballs/m ²	Socio-economic Impacts to Shoreline Users/Risk Factor 4C-1 and 2
Oil Slick/Tarballs	Brown to Black	0.1 mm	100 g/m ²	~12-14 tarballs/m ²	Ecological Impacts to Shoreline Habitats/Risk Factor 3C-1 and 2

Potential Impacts to the Water Column

Impacts to the water column from an oil release from the *Esso Gettysburg* will be determined by the volume of leakage. Because oil from sunken vessels will be released at low pressures, the droplet sizes will be large enough for the oil to float to the surface. Therefore, impacts to water column resources will result from the natural dispersion of the floating oil slicks on the surface, which is limited to about the top 33 feet. The metric used for ranking impacts to the water column is the area of water surface in mi² that has been contaminated by 1 part per billion (ppb) oil to a depth of 33 feet. At 1 ppb, there are likely to be impacts to sensitive organisms in the water column and potential tainting of seafood, so this concentration is used as a screening threshold for both the ecological and socio-economic risk factors for water column resource impacts. To assist planners in understanding the scale of potential impacts for different leakage volumes, a regression curve was generated for the water column volume oiled using the five volume scenarios, which is shown in Figure 2-1. Using this figure, the water column impacts can be estimated for any spill volume.

² French, D., M. Reed, K. Jayko, S. Feng, H. Rines, S. Pavignano, T. Isaji, S. Puckett, A. Keller, F. W. French III, D. Gifford, J. McCue, G. Brown, E. MacDonald, J. Quirk, S. Natzke, R. Bishop, M. Welsh, M. Phillips and B.S. Ingram, 1996. The CERCLA type A natural resource damage assessment model for coastal and marine environments (NRDAM/CME), Technical Documentation, Vol. I - V. Office of Environmental Policy and Compliance, U.S. Dept. of the Interior, Washington, DC.




Figure 2-1: Regression curve for estimating the volume of water column impacted as a function of spill volume for the *Esso Gettysburg*.

Potential Water Surface Slick

The slick size from an oil release from the *Esso Gettysburg* is a function of the quantity released. The estimated water surface coverage by a fresh slick (the total water surface area “swept” by oil over time) for the various scenarios is shown in Table 2-3, as the mean result of the 200 model runs. Note that this is an estimate of total water surface affected over a 30-day period. In the model, the representative crude oil used for this analysis spreads to a minimum thickness of approximately 975 g/m^2 , and is not able to spread any thinner. As a result, water surface oiling results are identical for the 0.01 g/m^2 and 10 g/m^2 thresholds. The slick will not be continuous but rather be broken and patchy due to the subsurface release of the oil. Surface expression is likely to be in the form of sheens, tarballs, and streamers.

Table 2-3: Estimated slick area swept on water for oil release scenarios from the *Esso Gettysburg*.

Scenario Type	Oil Volume (bbl)	Estimated Slick Area Swept Mean of All Models	
		0.01 g/m^2	10 g/m^2
Chronic	132	13,700 mi^2	13,600 mi^2
Episodic	1,320	44,500 mi^2	44,200 mi^2
Most Probable	13,200	152,000 mi^2	151,000 mi^2
Large	66,000	382,000 mi^2	380,000 mi^2
Worst Case Discharge	132,000	597,000 mi^2	595,000 mi^2

The location, size, shape, and spread of the oil slick(s) from an oil release will depend on environmental conditions, including winds and currents, at the time of release and in its aftermath. The areas potentially affected by oil slicks, given that we cannot predict when the spill might occur and the range of possible wind and current conditions that might prevail after a release, are shown in Figure 2-2 and Figure 2-3 using the Most Probable volume and the socio-economic and ecological thresholds.




Figure 2-2: Probability of surface oil (exceeding 0.01 g/m²) from the Most Probable spill of 13,200 bbl of medium crude oil from the *Esso Gettysburg* at the threshold for socio-economic resources at risk.




Figure 2-3: Probability of surface oil (exceeding 10 g/m²) from the Most Probable spill of 13,200 bbl of medium crude oil from the *Esso Gettysburg* at the threshold for ecological resources at risk.

The maximum potential cumulative area swept by oil slicks at some time after a Most Probable Discharge is shown in Figure 2-4 as the timing of oil movements.




Figure 2-4: Water surface oiling from the Most Probable spill of 13,200 bbl of medium crude oil from the Esso Gettysburg shown as the area over which the oil spreads at different time intervals.

The actual area affected by a release will be determined by the volume of leakage, whether it is from one or more tanks at a time. To assist planners in understanding the scale of potential impacts for different leakage volumes, a regression curve was generated for the water surface area oiled using the five volume scenarios, which is shown in Figure 2-5. Using this figure, the area of water surface with a barely visible sheen can be estimated for any spill volume.




Figure 2-5: Regression curve for estimating the amount of water surface oiling as a function of spill volume for the *Esso Gettysburg*, showing both the ecological threshold of 10 g/m² and socio-economic threshold of 0.01 g/m². The curves are so similar that they plot on top of each other.

Potential Shoreline Impacts

Based on these modeling results, shorelines from Cape Lookout, North Carolina to the entrance to Chesapeake Bay are at risk. Figure 2-6 shows the probability of oil stranding on the shoreline at concentrations that exceed the threshold of 1 g/m², for the Most Probable release of 13,200 bbl. However, the specific areas that would be oiled will depend on the currents and winds at the time of the oil release(s), as well as on the amount of oil released. Figure 2-7 shows the single oil spill scenario that resulted in the maximum extent of shoreline oiling for the Most Probable volume. Estimated miles of shoreline oiling above the threshold of 1 g/m² by scenario type are shown in Table 2-4.

Table 2-4: Estimated shoreline oiling from leakage from the *Esso Gettysburg*.

Scenario Type	Volume (bbl)	Estimated Miles of Shoreline Oiling Above 1 g/m ²			
		Rock/Gravel/Artificial	Sand	Wetland/Mudflat	Total
Chronic	132	1	8	0	9
Episodic	1,320	1	28	1	30
Most Probable	13,200	1	32	8	40
Large	66,000	1	36	12	50
Worst Case Discharge	132,000	2	40	15	57

The actual shore length affected by a release will be determined by the volume of leakage and environmental conditions during an actual release. To assist planners in scaling the potential impact for different leakage volumes, a regression curve was generated for the total shoreline length oiled using the five volume scenarios, which is shown in Figure 2-8. Using this figure, the shore length oiled can be estimated for any spill volume.




Figure 2-6: Probability of shoreline oiling (exceeding 1 g/m²) from the Most Probable Discharge of 13,200 bbl of medium crude oil from the *Esso Gettysburg*.




Figure 2-7: The extent and degree of shoreline oiling from the single model run of the Most Probable Discharge of 13,200 bbl of medium crude oil from the *Esso Gettysburg* that resulted in the greatest shoreline oiling.




Figure 2-8: Regression curve for estimating the amount of shoreline oiling at different thresholds as a function of spill volume for the *Esso Gettysburg*.

The worst case scenario for shoreline exposure along the potentially impacted area for the WCD volume (Table 2-5) and the Most Probable volume (Table 2-6) consists primarily of sand beaches. Salt marshes and tidal flats near tidal inlets are also at risk.

Table 2-5: Worst case scenario shoreline impact by habitat type and oil thickness for a leakage of 132,000 bbl from the *Esso Gettysburg*.

Shoreline/Habitat Type	Lighter Oiling Oil Thickness <1 mm Oil Thickness >1 g/m ²	Heavier Oiling Oil Thickness >1 mm Oil Thickness >100 g/m ²
Rocky and artificial shores/Gravel beaches	13 miles	3 miles
Sand beaches	170 miles	101 miles
Salt marshes and tidal flats	64 miles	6 miles

Table 2-6: Worst case scenario shoreline impact by habitat type and oil thickness for a leakage of 13,200 bbl from the *Esso Gettysburg*.

Shoreline/Habitat Type	Lighter Oiling Oil Thickness <1 mm Oil Thickness >1 g/m ²	Heavier Oiling Oil Thickness >1 mm Oil Thickness >100 g/m ²
Rocky and artificial shores/Gravel beaches	5 miles	2 miles
Sand beaches	135 miles	32 miles
Salt marshes and tidal flats	37 miles	0 miles

SECTION 3: ECOLOGICAL RESOURCES AT RISK

Ecological resources at risk from a catastrophic release of oil from the *Esso Gettysburg* include numerous guilds of birds (Table 3-1), particularly those sensitive to surface oiling while rafting or plunge diving to feed, that are present in nearshore/offshore waters. In addition, this region is important for nesting loggerhead sea turtles, migrating marine mammals, and commercially important fish and invertebrates, including some sensitive hardbottom habitats used by these species.

Table 3-1: Ecological resources at risk from a release of oil from the *Esso Gettysburg*.

FT = Federal threatened; FE = Federal endangered; ST = State threatened; SE = State endangered).

Species Group	Species Subgroup and Geography	Seasonal Presence
Seabirds	Seabirds <ul style="list-style-type: none"> Outer Continental Shelf offshore of Cape Hatteras, NC: greatest diversity of seabirds in SE U.S.; greatest density of tropical seabirds in SE U.S. Species include: shearwaters, storm petrels, Bermuda petrels Significant percentage of the global population of black-capped petrels (FE) may be present in <i>Sargassum</i> mats formed off Cape Hatteras and Gulf Stream off SE U.S. coast Audubon's shearwaters (50-75% of population) concentrate along the Continental Shelf edge off NC, extending northward to the VA border (~3,800 pairs) 	OCS: Ranges by species but Mar-Nov peak Petrels off NC/VA coast during summer through early fall and off SE U.S. coast in winter Shearwaters off of NC/VA: late summer
Pelagic Birds, Waterfowl, and Diving Birds	Coastal pelagic birds, waterfowl, diving birds <ul style="list-style-type: none"> Mouth of Chesapeake Bay has high concentrations of species that are abundant over shoals (e.g., loons, pelicans, cormorants, sea ducks, gulls, terns, alcids); scoters are 10X more abundant than other shoal species Outer Banks, inshore waters NC to VA: Key foraging area for gulls and terns; key migration corridor for loons and sea ducks; NC's largest population of northern gannet and red-breasted merganser Southeastern U.S. inshore/offshore waters: 150K loons, >15K pelicans, thousands of waterfowl, 100K of cormorants and terns, millions of gulls 	Winter use of shoals (Dec-Mar); Summer use of shoals likely farther north Spring/summer, for terns, gulls; Spring/fall for loons, sea ducks; winter for waterfowl, gannets and red-breasted mergansers
Sea Ducks	Sea ducks (includes mean and max distance of flocks to shore, 2009-2010 data) off North Carolina <ul style="list-style-type: none"> Surf scoter - 2 nm/8 nm: 0-41K Black scoter -2 nm/13 nm: 3.5-13K Bufflehead, mergansers, goldeneyes (<1 nm/7-14 nm): 12K 	Sea ducks surveyed in winter (peak abundances); Migration from fall to spring (Oct-Apr)
Shorebirds and Colonial Nesting Birds	<ul style="list-style-type: none"> Outer Banks, Cape Hatteras, and Cape Lookout: Globally important for coastal birds with 365+ species Battery and Bald Head Islands, NC: Largest colonies of wading birds in NC; globally significant site with >10K nesting pairs of white ibises 	Winter migration stop for plovers; Colonial and beach nesters peak Apr-Aug Wading and shorebirds typically present year round
Sea Turtles	Nesting (annual counts, by state, along shorelines with most probable impacts). NC nesting <ul style="list-style-type: none"> 650+ Loggerhead (FT) <20 Green (FT) <10 Leatherback (FE) Distribution: <ul style="list-style-type: none"> Offshore hot spots not well known Young associated with <i>Sargassum</i> mats off Cape Hatteras Bays and sounds are foraging grounds for juvenile green, loggerhead, and Kemp's ridley (FE) 	Nesting season: Loggerheads/Greens (NC) Adults: May-Aug Hatching: Jul-Oct Leatherbacks (NC) Adults: Mar-Jul Hatching: May-Oct In water: Year round with Apr-Dec peak

Species Group	Species Subgroup and Geography	Seasonal Presence
Marine Mammals	<p><i>Baleen whales</i>: Primarily North Atlantic right whale (FE) with occasional humpback whale (FE), and minke whales</p> <ul style="list-style-type: none"> Right whales are critically endangered (<400 individuals left); coastal waters are used as calving grounds <p><i>Inshore cetaceans</i>: Bottlenose dolphins frequently use coastal waters including creeks, bays, and sounds throughout potential spill area.</p> <p><i>Offshore cetaceans</i>: Risso's dolphin, striped dolphin, clymene dolphin, Atlantic spotted dolphin, spinner dolphin, short-finned pilot whales, pantropical spotted dolphin; Often associated with shelf edge features, convergence zones (fronts), and <i>Sargassum</i> mats (summer)</p> <p><i>Pinnipeds and Sirenians</i>:</p> <ul style="list-style-type: none"> Juvenile harbor and hooded seals can sometimes occur as far south as N. FL during the winter West Indian manatees common summer sightings as far north as NC 	<p>Adults migrate from feeding grounds in North Atlantic to breeding grounds further south in the winter; Right whales with calf Nov-Mar</p> <p>Bottlenose dolphins present year round</p> <p>Harbor and hooded seals present during winter</p> <p>Manatees year round and coastal waters during summer</p>
Fish and Inverts	<p>Coastal ocean waters support many valuable fisheries and/or species of concern in the region:</p> <ul style="list-style-type: none"> <i>Benthic or bottom associated</i>: Snapper, grouper, black sea bass, butter fish, goose fish, shrimp (white, pink, brown, and rock), golden crab, and other reef species <i>Midwater</i>: Atlantic mackerel, Spanish mackerel, shortfin squid, bluefish, menhaden, cero, cobia <i>Pelagic</i>: Bluefin tuna, yellowfin tuna, wahoo, dolphinfish, bigeye tuna, swordfish, marlin, sailfish <i>Diadromous</i>: Alewife, blueback herring, American shad, hickory shad, Atlantic tomcod, American eel, Atlantic sturgeon (Fed. species of concern), shortnose sturgeon (FE), and striped bass <i>Estuarine dependent</i>: Southern flounder, redfish, spotted seatrout, blue crab, atlantic croaker, spot, weakfish, shrimp <i>Estuarine resident</i>: Eastern oyster <p>Important concentration/conservation areas are:</p> <ul style="list-style-type: none"> Pelagic species can be more concentrated around the shelf break and at oceanographic fronts in the region The Point (offshore of Cape Hatteras) – Essential Fish Habitat/Habitats Areas of Particular concern (EFH/HAPC) for coastal migratory pelagics and dolphin/wahoo Ten Fathom Ledge – South of Cape Lookout Big Rock – SE of Cape Lookout <i>Sargassum</i> off Cape Hatteras, NC is important habitat for juvenile of some pelagic fish species (i.e., dolphinfish, jacks, triggerfish) 	<p>Benthic and midwater species are present throughout the year</p> <p>Bluefin tuna present fall-spring with other pelagic fish present year round</p> <p>Anadromous fish migrate inshore to spawn in fresh water in spring</p> <p>American eel migrates offshore to spawn in winter</p> <p>Estuarine dependent fish migrate offshore in fall/winter to spawn; Juveniles and adults use estuaries in spring/summer</p>
Benthic Habitats	<p>Submerged aquatic vegetation is critical to numerous species and occurs inside of bays and sounds throughout the region</p> <p>Scattered hard-bottom sites are located off NC and are considered HAPC for reef-associated fishes (including the areas listed above)</p>	<p>Year round</p>

The Environmental Sensitivity Index (ESI) atlases for the potentially impacted coastal areas from a leak from the *Esso Gettysburg* are generally available at each U.S. Coast Guard Sector. They can also be downloaded at: <http://response.restoration.noaa.gov/esi>. These maps show detailed spatial information on

the distribution of sensitive shoreline habitats, biological resources, and human-use resources. The tables on the back of the maps provide more detailed life-history information for each species and location. The ESI atlases should be consulted to assess the potential environmental resources at risk for specific spill scenarios. In addition, the Geographic Response Plans within the Area Contingency Plans prepared by the Area Committee for each U.S. Coast Guard Sector have detailed information on the nearshore and shoreline ecological resources at risk and should be consulted.

Ecological Risk Factors

Risk Factor 3: Impacts to Ecological Resources at Risk (EcoRAR)

Ecological resources include plants and animals (e.g., fish, birds, invertebrates, and mammals), as well as the habitats in which they live. All impact factors are based on a Worst Case and the Most Probable Discharge oil release from the wreck. Risk factors for ecological resources at risk (EcoRAR) are divided into three categories:

- Impacts to the water column and resources in the water column;
- Impacts to the water surface and resources on the water surface; and
- Impacts to the shoreline and resources on the shoreline.

The impacts from an oil release from the wreck would depend greatly on the direction in which the oil slick moves, which would, in turn, depend on wind direction and currents at the time of and after the oil release. Impacts are characterized in the risk analysis based on the likelihood of any measurable impact, as well as the degree of impact that would be expected if there is an impact. The measure of the degree of impact is based on the median case for which there is at least some impact. The median case is the “middle case” – half of the cases with significant impacts have less impact than this case, and half have more.

For each of the three ecological resources at risk categories, risk is defined as:

- The **probability of oiling** over a certain threshold (i.e., the likelihood that there will be an impact to ecological resources over a certain minimal amount); and
- The **degree of oiling** (the magnitude or amount of that impact).

As a reminder, the ecological impact thresholds are: 1 ppb aromatics for water column impacts; 10 g/m² for water surface impacts; and 100 g/m² for shoreline impacts.

In the following sections, the definition of low, medium, and high for each ecological risk factor is provided. Also, the classification for the *Esso Gettysburg* is provided, both as text and as **shading** of the applicable degree of risk bullet, for the WCD release of 132,000 bbl and **a border** around the Most Probable Discharge of 13,200 bbl.

Risk Factor 3A: Water Column Impacts to EcoRAR

Water column impacts occur beneath the water surface. The ecological resources at risk for water column impacts are fish, marine mammals, and invertebrates (e.g., shellfish, and small organisms that are food for

larger organisms in the food chain). These organisms can be affected by toxic components in the oil. The threshold for water column impact to ecological resources at risk is a dissolved aromatic hydrocarbons concentration of 1 ppb (i.e., 1 part total dissolved aromatics per one billion parts water). Dissolved aromatic hydrocarbons are the most toxic part of the oil. At this concentration and above, one would expect impacts to organisms in the water column.

Risk Factor 3A-1: Water Column Probability of Oiling of EcoRAR

This risk factor reflects the probability that at least 0.2 mi² of the upper 33 feet of the water column would be contaminated with a high enough concentration of oil to cause ecological impacts. The three risk scores for water column oiling probability are:

- **Low Oiling Probability:** Probability = <10%
- **Medium Oiling Probability:** Probability = 10 – 50%
- **High Oiling Probability:** Probability > 50%

Risk Factor 3A-2: Water Column Degree of Oiling of EcoRAR

The degree of oiling of the water column reflects the total volume of water that would be contaminated by oil at a concentration high enough to cause impacts. The three categories of impact are:

- **Low Impact:** impact on less than 0.2 mi² of the upper 33 feet of the water column at the threshold level
- **Medium Impact:** impact on 0.2 to 200 mi² of the upper 33 feet of the water column at the threshold level
- **High Impact:** impact on more than 200 mi² of the upper 33 feet of the water column at the threshold level

The *Esso Gettysburg* is classified as High Risk for oiling probability for water column ecological resources for the WCD of 132,000 bbl because 72% of the model runs resulted in contamination of more than 0.2 mi² of the upper 33 feet of the water column above the threshold of 1 ppb aromatics. It is classified as High Risk for degree of oiling because the mean volume of water contaminated was 341 mi² of the upper 33 feet of the water column. For the Most Probable Discharge of 13,200 bbl, the *Esso Gettysburg* is classified as High Risk for oiling probability for water column ecological resources because 100% of the model runs resulted in contamination of more than 0.2 mi² of the upper 33 feet of the water column above the threshold of 1 ppb aromatics. It is classified as Medium Risk for degree of oiling because the mean volume of water contaminated was 81 mi² of the upper 33 feet of the water column.

Risk Factor 3B: Water Surface Impacts to EcoRAR

Ecological resources at risk at the water surface include surface feeding and diving sea birds, sea turtles, and marine mammals. These organisms can be affected by the toxicity of the oil as well as from coating with oil. The threshold for water surface oiling impact to ecological resources at risk is 10 g/m² (10 grams of floating oil per square meter of water surface). At this concentration and above, one would expect impacts to birds and other animals that spend time on the water surface.

Risk Factor 3B-1: Water Surface Probability of Oiling of EcoRAR

This risk factor reflects the probability that at least 1,000 mi² of the water surface would be affected by enough oil to cause impacts to ecological resources. The three risk scores for oiling are:

- **Low Oiling Probability:** Probability = <10%

- **Medium Oiling Probability:** Probability = 10 – 50%

- **High Oiling Probability:** Probability > 50%

Risk Factor 3B-2: Water Surface Degree of Oiling of EcoRAR

The degree of oiling of the water surface reflects the total amount of oil that would affect the water surface in the event of a discharge from the vessel. The three categories of impact are:

- **Low Impact:** less than 1,000 mi² of water surface impact at the threshold level
- **Medium Impact:** 1,000 to 10,000 mi² of water surface impact at the threshold level
- **High Impact:** more than 10,000 mi² of water surface impact at the threshold level

The *Esso Gettysburg* is classified as High Risk for oiling probability for water surface ecological resources for the WCD because 100% of the model runs resulted in at least 1,000 mi² of the water surface affected above the threshold of 10 g/m². It is classified as High Risk for degree of oiling because the mean area of water contaminated was 595,000 mi². It is classified as High Risk for oiling probability for water surface ecological resources for the Most Probable Discharge because 100% of the model runs resulted in at least 1,000 mi² of the water surface affected above the threshold of 10 g/m². It is classified as High Risk for degree of oiling because the mean area of water contaminated was 151,000 mi².

Risk Factor 3C: Shoreline Impacts to EcoRAR

The impacts to different types of shorelines vary based on their type and the organisms that live on them. In this risk analysis, shorelines have been weighted by their degree of sensitivity to oiling. Wetlands are the most sensitive (weighted as “3” in the impact modeling), rocky and gravel shores are moderately sensitive (weighted as “2”), and sand beaches (weighted as “1”) are the least sensitive to ecological impacts of oil.

Risk Factor 3C-1: Shoreline Probability of Oiling of EcoRAR

This risk factor reflects the probability that the shoreline would be coated by enough oil to cause impacts to shoreline organisms. The threshold for shoreline oiling impacts to ecological resources at risk is 100 g/m² (i.e., 100 grams of oil per square meter of shoreline). The three risk scores for oiling are:

- **Low Oiling Probability:** Probability = <10%
- **Medium Oiling Probability:** Probability = 10 – 50%
- **High Oiling Probability:** Probability > 50%

Risk Factor 3C-2: Shoreline Degree of Oiling of EcoRAR

The degree of oiling of the shoreline reflects the length of shorelines oiled by at least 100 g/m² in the event of a discharge from the vessel. The three categories of impact are:

- **Low Impact:** less than 10 miles of shoreline impacted at the threshold level
- **Medium Impact:** 10 - 100 miles of shoreline impacted at the threshold level
- **High Impact:** more than 100 miles of shoreline impacted at the threshold level

The *Esso Gettysburg* is classified as Medium Risk for oiling probability for shoreline ecological resources for the WCD because 12% of the model runs resulted in shorelines affected above the threshold of 100 g/m². It is classified as Medium Risk for degree of oiling because the mean weighted length of shoreline

contaminated was 37 miles. The *Esso Gettysburg* is classified as Low Risk for oiling probability to shoreline ecological resources for the Most Probable Discharge because 6% of the model runs resulted in shorelines affected above the threshold of 100 g/m². It is classified as Medium Risk for degree of oiling because the mean weighted length of shoreline contaminated was 13 miles.

Considering the modeled risk scores and the ecological resources at risk, the ecological risk from potential releases of the WCD of 132,000 bbl of crude oil from the *Esso Gettysburg* is summarized as listed below and indicated in the far-right column in Table 3-2:

- Water column resources – Medium, because the area of highest exposure occurs in open shelf waters without any known concentrations of sensitive upper water column resources
- Water surface resources – High, because of the seasonally very large number wintering, nesting, and migratory birds that use ocean, coastal, and estuarine habitats at risk and importance of the area to adult and juvenile sea turtles. It should be noted that oil on the surface will not be continuous but rather be broken and patchy and in the form of sheens, tarballs, and streamers
- Shoreline resources – Medium, because any spilled oil is likely to undergo extensive weathering at sea prior to stranding mostly on sand beaches, although marshes and tidal flats are also at risk

Table 3-2: Ecological risk factor scores for the **Worst Case Discharge of 132,000 bbl** of crude oil from the *Esso Gettysburg*.

Risk Factor	Risk Score			Explanation of Risk Score	Final Score
3A-1: Water Column Probability EcoRAR Oiling	Low	Medium	High	72% of the model runs resulted in at least 0.2 mi ² of the upper 33 feet of the water column contaminated above 1 ppb aromatics	Med
3A-2: Water Column Degree EcoRAR Oiling	Low	Medium	High	The mean volume of water contaminated above 1 ppb was 340 mi ² of the upper 33 feet of the water column	
3B-1: Water Surface Probability EcoRAR Oiling	Low	Medium	High	100% of the model runs resulted in at least 1,000 mi ² of water surface covered by at least 10 g/m ²	High
3B-2: Water Surface Degree EcoRAR Oiling	Low	Medium	High	The mean area of water contaminated above 10 g/m ² was 595,000 mi ²	
3C-1: Shoreline Probability EcoRAR Oiling	Low	Medium	High	12% of the model runs resulted in shoreline oiling of 100 g/m ²	Med
3C-2: Shoreline Degree EcoRAR Oiling	Low	Medium	High	The length of shoreline contaminated by at least 100 g/m ² was 37 mi	

For the Most Probable Discharge of 13,200 bbl, the ecological risk from potential releases of crude oil from the *Esso Gettysburg* is summarized as listed below and indicated in the far-right column in Table 3-3:

- Water column resources – Medium, because the moderate-sized area of highest exposure occurs in open shelf waters without any known concentrations of sensitive upper water column resources
- Water surface resources – High, because of the seasonally very large number wintering, nesting, and migratory birds that use ocean, coastal, and estuarine habitats at risk and importance of the area to adult and juvenile sea turtles. It should be noted that oil on the surface will not be continuous but rather be broken and patchy and in the form of sheens, tarballs, and streamers
- Shoreline resources – Low, because fewer miles of shoreline, mostly sand beaches, are at risk from oil that is likely to undergo extensive weathering at sea prior to stranding

Table 3-3: Ecological risk factor scores for the **Most Probable Discharge of 13,200 bbl** of crude oil from the *Esso Gettysburg*.

Risk Factor	Risk Score			Explanation of Risk Score	Final Score
3A-1: Water Column Probability EcoRAR Oiling	Low	Medium	High	100% of the model runs resulted in at least 0.2 mi ² of the upper 33 feet of the water column contaminated above 1 ppb aromatics	Med
3A-2: Water Column Degree EcoRAR Oiling	Low	Medium	High	The mean volume of water contaminated above 1 ppb was 81 mi ² of the upper 33 feet of the water column	
3B-1: Water Surface Probability EcoRAR Oiling	Low	Medium	High	100% of the model runs resulted in at least 1,000 mi ² of water surface covered by at least 10 g/m ²	High
3B-2: Water Surface Degree EcoRAR Oiling	Low	Medium	High	The mean area of water contaminated above 10 g/m ² was 151,000 mi ²	
3C-1: Shoreline Probability EcoRAR Oiling	Low	Medium	High	6% of the model runs resulted in shoreline oiling of 100 g/m ²	Low
3C-2: Shoreline Degree EcoRAR Oiling	Low	Medium	High	The length of shoreline contaminated by at least 100 g/m ² was 13 mi	

SECTION 4: SOCIO-ECONOMIC RESOURCES AT RISK

In addition to natural resource impacts, spills from sunken wrecks have the potential to cause significant social and economic impacts. Socio-economic resources potentially at risk from oiling are listed in Table 4-1 and shown in Figures 4-1 and 4-2. The potential economic impacts include disruption of coastal economic activities such as commercial and recreational fishing, boating, vacationing, commercial shipping, and other activities that may become claims following a spill.

Socio-economic resources in the areas potentially affected by a release from the *George MacDonald* include very highly utilized recreational beaches from Maryland to northeastern Florida during summer, but also during spring and fall for shore fishing. Four national seashores and two coastal national monuments would potentially be affected. Many areas along the entire potential spill zone are widely popular seaside resorts and support recreational activities such as boating, diving, sightseeing, sailing, fishing, and wildlife viewing. The Gray's Reef National Marine Sanctuary off Georgia would also potentially be affected, along with a large number of coastal state parks.

There are shipping lanes to several ports that could be impacted by a release with a total of nearly 9,000 annual port calls annually with a total of over 382 million tonnage. Commercial fishing is economically important to the region. Regional commercial landings for 2010 exceeded \$137 million with fishing fleets from Maryland to Florida potentially impacted by a release.

In addition to the ESI Atlases, the Geographic Response Plans within the Area Contingency Plans prepared by the Area Committee for each U.S. Coast Guard Sector have detailed information on important socio-economic resources at risk and should be consulted.

Spill response costs for a release of oil from the *Esso Gettysburg* would be dependent on volume of oil released and specific areas impacted. The specific shoreline impacts and spread of the oil would determine the response required and the costs for that response.

Table 4-1: Socio-economic resources at risk from a release of oil from the *Esso Gettysburg*.

Resource Type	Resource Name	Economic Activities
Tourist Beaches	Myrtle Beach, SC	Potentially affected beach resorts and beach-front communities in North Carolina and South Carolina provide recreational activities (e.g., swimming, boating, recreational fishing, wildlife viewing, nature study, sports, dining, camping, and amusement parks) with substantial income for local communities and state tax income. Much of the coast is lined with economically valuable beach resorts and residential communities, with recreational activities concentrated in late spring to early fall months.
National Marine Sanctuary	Gray's Reef National Marine Sanctuary (GA)	Gray's Reef National Marine Sanctuary is one of the largest near shore live-bottom reefs in the southeastern U.S. The Sanctuary is popular with recreational anglers, boaters, and more experienced divers.
National Wildlife Refuges	Wallops Island NWR (VA) Eastern Shore of Virginia NWR (VA)	National wildlife refuges in five states may be impacted. These federally-managed and protected lands provide

Resource Type	Resource Name	Economic Activities
	Back Bay NWR (VA) Mackay Island NWR (NC) Currituck NWR (NC) Pea Island NWR (NC) Cedar Island NWR (NC) Waccamaw NWR (SC) Cape Romain NWR (SC) Ernest F. Hollings ACE Basin NWR (SC) Pickney Island NWR (SC) Savannah NWR (SC) Tybee NWR (SC)	refuges and conservation areas for sensitive species and habitats.
National Seashores	Cape Hatteras National Seashore, NC Assateague Isl. National Seashore, MD/VA	National seashores provide recreation for local and tourist populations as well as preserve and protect the nation's natural shoreline treasures. National seashores are coastal areas federally designated as being of natural and recreational significance as a preserved area. Cape Hatteras is known for its Bodie Island and Cape Hatteras Lighthouses. Popular recreation activities include windsurfing, birdwatching, fishing, shell collecting, and kayaking. The barrier islands provide refuge for the endangered piping plover, seabeach amaranth, and sea turtles.
State Parks	Myrtle Beach SP, SC Huntington Beach SP, SC	Coastal state parks are significant recreational resources for the public (e.g., swimming, boating, recreational fishing, wildlife viewing, nature study, sports, dining, camping, and amusement parks). They provide income to the states. State parks in several states are potentially impacted. Many of these recreational activities are limited to or concentrated into the late spring into early fall months.
Commercial Fishing	A number of fishing fleets use potentially affected waters for commercial fishing.	
	Chincoteague, Virginia	Total Landings (2010): \$3.5M
	Ocean City, Maryland	Total Landings (2010): \$8.8M
	Chincoteague, Virginia	Total Landings (2010): \$3.5M
	Beaufort-Morehead City, NC	Total Landings (2010): \$9.2M
	Belhaven-Washington, NC	Total Landings (2010): \$3.7M
	Elizabeth City, NC	Total Landings (2010): \$5.4M
	Engelhard-Swanquarter, NC	Total Landings (2010): \$10.6M
	Oriental-Vandemere, NC	Total Landings (2010): \$8.4M
	Sneads Ferry-Swansboro, NC	Total Landings (2010): \$5.4M
	Wanchese-Stumpy Point, NC	Total Landings (2010): \$22.0M
	Brunswick, GA	Total Landings (2010): \$5.1M
	Cape Canaveral, FL	Total Landings (2010): \$6.5M
	Charleston-Mt. Pleasant, SC	Total Landings (2010): \$9.9M
	Darien-Bellville, GA	Total Landings (2010): \$5.2M
	Fernandina Beach, FL	Total Landings (2010): \$4.7M
	Georgetown, SC	Total Landings (2010): \$6.0M
	Mayport, FL	Total Landings (2010): \$11.0M
	Savannah, GA	Total Landings (2010): \$5.0M
	Thunderbolt, GA	Total Landings (2010): \$3.4M
Ports	There are a number of significant commercial ports along the Atlantic coast that could potentially be impacted by spillage and spill response activities. The port call numbers below are for large vessels only. There are many more, smaller vessels (under 400 GRT) that also use these ports.	

Resource Type	Resource Name	Economic Activities
	Baltimore, MD	2,100 port calls annually
	Morehead City, NC	85 port calls annually
	Wilmington, NC	550 port calls annually
	Brunswick, GA	304 port calls annually
	Charleston, SC	1,818 port calls annually
	Elba Is., GA	37 port calls annually
	Fernandina, FL	3 port calls annually
	Jacksonville, FL	1,641 port calls annually
	Port Canaveral, FL	38 port calls annually
	Savannah, GA	2,406 port calls annually




Figure 4-1: Tribal lands, ports and commercial fishing fleets at risk from a release from the *Esso Gettysburg*. (Note that there are no tribal lands at risk.)




Figure 4-2: Beaches, coastal state parks, national marine sanctuary, and Federal protected areas at risk from a release from the *Esso Gettysburg*.

Socio-Economic Risk Factors

Risk Factor 4: Impacts to Socio-economic Resources at Risk (SRAR)

Socio-economic resources at risk (SRAR) include potentially impacted resources that have some economic value, including commercial and recreational fishing, tourist beaches, private property, etc. All impact factors are evaluated for both the Worst Case and the Most Probable Discharge oil release from the wreck. Risk factors for socio-economic resources at risk are divided into three categories:

- **Water Column:** Impacts to the water column and to socio-economic resources in the water column (i.e., fish and invertebrates that have economic value);
- **Water Surface:** Impacts to the water surface and resources on the water surface (i.e., boating and commercial fishing); and
- **Shoreline:** Impacts to the shoreline and resources on the shoreline (i.e., beaches, real property).

The impacts from an oil release from the wreck would depend greatly on the direction in which the oil slick moves, which would, in turn, depend on wind direction and currents at the time of and after the oil release. Impacts are characterized in the risk analysis based on the likelihood of any measurable impact, as well as the degree of impact that would be expected if there were one. The measure of the degree of impact is based on the median case for which there is at least some impact. The median case is the “middle case” – half of the cases with significant impacts have less impact than this case, and half have more.

For each of the three socio-economic resources at risk categories, risk is classified with regard to:

- The **probability of oiling** over a certain threshold (i.e., the likelihood that there will be exposure to socio-economic resources over a certain minimal amount known to cause impacts); and
- The **degree of oiling** (the magnitude or amount of that exposure over the threshold known to cause impacts).

As a reminder, the socio-economic impact thresholds are: 1 ppb aromatics for water column impacts; 0.01 g/m² for water surface impacts; and 1 g/m² for shoreline impacts.

In the following sections, the definition of low, medium, and high for each socio-economic risk factor is provided. Also, in the text classification for the *Esso Gettysburg* shading indicates the degree of risk for the WCD release of 132,000 bbl and a border indicates degree of risk for the Most Probable Discharge of 13,200 bbl of crude oil.

Risk Factor 4A-1: Water Column: Probability of Oiling of SRAR

This risk factor reflects the probability that at least 0.2 mi² of the upper 33 feet of the water column would be contaminated with a high enough concentration of oil to cause socio-economic impacts. The threshold for water column impact to socio-economic resources at risk is an oil concentration of 1 ppb (i.e., 1 part oil per one billion parts water). At this concentration and above, one would expect impacts and potential tainting to socio-economic resources (e.g., fish and shellfish) in the water column; this concentration is used as a screening threshold for both the ecological and socio-economic risk factors.

The three risk scores for oiling are:

- **Low Oiling Probability:** Probability = <10%
- **Medium Oiling Probability:** Probability = 10 – 50%
- **High Oiling Probability:** Probability > 50%

Risk Factor 4A-2: Water Column Degree of Oiling of SRAR

The degree of oiling of the water column reflects the total amount of oil that would affect the water column in the event of a discharge from the vessel. The three categories of impact are:

- **Low Impact:** impact on less than 0.2 mi² of the upper 33 feet of the water column at the threshold level
- **Medium Impact:** impact on 0.2 to 200 mi² of the upper 33 feet of the water column at the threshold level
- **High Impact:** impact on more than 200 mi² of the upper 33 feet of the water column at the threshold level

The *Esso Gettysburg* is classified as High Risk for both oiling probability and degree of oiling for water column socio-economic resources for the WCD of 132,000 bbl because 72% of the model runs resulted in contamination of more than 0.2 mi² of the upper 33 feet of the water column above the threshold of 1 ppb aromatics, and the mean volume of water contaminated was 341 mi² of the upper 33 feet of the water column. For the Most Probable Discharge of 13,200 bbl, the *Esso Gettysburg* is classified as High Risk for oiling probability for water column socio-economic resources because 100% of the model runs resulted in contamination of more than 0.2 mi² of the upper 33 feet of the water column above the

threshold of 1 ppb aromatics. It is classified as Medium Risk for degree of oiling because the mean volume of water contaminated was 81 mi² of the upper 33 feet of the water column.

Risk Factor 4B-1: Water Surface Probability of Oiling of SRAR

This risk factor reflects the probability that at least 1,000 mi² of the water surface would be affected by enough oil to cause impacts to socio-economic resources. The three risk scores for oiling are:

- **Low Oiling Probability:** Probability = <10%
- **Medium Oiling Probability:** Probability = 10 – 50%
- **High Oiling Probability:** Probability > 50%

The threshold level for water surface impacts to socio-economic resources at risk is 0.01 g/m² (i.e., 0.01 grams of floating oil per square meter of water surface). At this concentration and above, one would expect impacts to socio-economic resources on the water surface.

Risk Factor 4B-2: Water Surface Degree of Oiling of SRAR

The degree of oiling of the water surface reflects the total amount of oil that would affect the water surface in the event of a discharge from the vessel. The three categories of impact are:

- **Low Impact:** less than 1,000 mi² of water surface impact at the threshold level
- **Medium Impact:** 1,000 to 10,000 mi² of water surface impact at the threshold level
- **High Impact:** more than 10,000 mi² of water surface impact at the threshold level

The *Esso Gettysburg* is classified as High Risk for both oiling probability and degree of oiling for water surface socio-economic resources for the WCD because 100% of the model runs resulted in at least 1,000 mi² of the water surface affected above the threshold of 0.01 g/m², and the mean area of water contaminated was 597,000 mi². The *Esso Gettysburg* is classified as High Risk for both oiling probability and degree of oiling for water surface socio-economic resources for the Most Probable Discharge because 100% of the model runs resulted in at least 1,000 mi² of the water surface affected above the threshold of 0.01 g/m², and the mean area of water contaminated was 152,400 mi².

Risk Factor 4C: Shoreline Impacts to SRAR

The impacts to different types of shorelines vary based on economic value. In this risk analysis, shorelines have been weighted by their degree of sensitivity to oiling. Sand beaches are the most economically valued shorelines (weighted as “3” in the impact analysis), rocky and gravel shores are moderately valued (weighted as “2”), and wetlands are the least economically valued shorelines (weighted as “1”). Note that these values differ from the ecological values of these three shoreline types.

Risk Factor 4C-1: Shoreline Probability of Oiling of SRAR

This risk factor reflects the probability that the shoreline would be coated by enough oil to cause impacts to shoreline users. The threshold for impacts to shoreline SRAR is 1 g/m² (i.e., 1 gram of oil per square meter of shoreline). The three risk scores for oiling are:

- **Low Oiling Probability:** Probability = <10%
- **Medium Oiling Probability:** Probability = 10 – 50%
- **High Oiling Probability:** Probability > 50%

Risk Factor 4C-2: Shoreline Degree of Oiling of SRAR

The degree of oiling of the shoreline reflects the total amount of oil that would affect the shoreline in the event of a discharge from the vessel. The three categories of impact are:

- **Low Impact:** less than 10 miles of shoreline impacted at threshold level
- **Medium Impact:** 10 - 100 miles of shoreline impacted at threshold level
- **High Impact:** more than 100 miles of shoreline impacted at threshold level

The *Esso Gettysburg* is classified as Medium Risk for oiling probability for shoreline socio-economic resources for the WCD because 12% of the model runs resulted in shorelines affected above the threshold of 1 g/m². It is classified as High Risk for degree of oiling because the mean length of weighted shoreline contaminated was 139 miles. The *Esso Gettysburg* is classified as Medium Risk for oiling probability and High Risk for degree of oiling for shoreline socio-economic resources for the Most Probable Discharge as 14% of the model runs resulted in shorelines affected above the threshold of 1 g/m², and the mean length of weighted shoreline contaminated was 104 miles.

Considering the modeled risk scores and the socio-economic resources at risk, the socio-economic risk from potential releases of the WCD of 132,000 bbl of crude oil from the *Esso Gettysburg* is summarized as listed below and indicated in the far-right column in Table 4-2:

- Water column resources – High, because a large area of water column would be impacted in areas with fishing grounds and other offshore resources (e.g., marine sanctuary)
- Water surface resources – High, because a large area of water surface would be impacted offshore in areas with shipping lanes and other offshore activities. It should be noted that oil on the surface will not be continuous but rather be broken and patchy and in the form of sheens, tarballs, and streamers
- Shoreline resources – High, because a relatively large length of shoreline would be impacted in areas with high-value and sensitive resources

Table 4-2: Socio-economic risk factor ranks for the **Worst Case Discharge of 132,000 bbl** of crude oil from the *Esso Gettysburg*.

Risk Factor	Risk Score			Explanation of Risk Score	Final Score
4A-1: Water Column Probability SRAR Oiling	Low	Medium	High	72% of the model runs resulted in at least 0.2 mi ² of the upper 33 feet of the water column contaminated above 1 ppb aromatics	High
4A-2: Water Column Degree SRAR Oiling	Low	Medium	High	The mean volume of water contaminated above 1 ppb was 341 mi ² of the upper 33 feet of the water column	
4B-1: Water Surface Probability SRAR Oiling	Low	Medium	High	100% of the model runs resulted in at least 1,000 mi ² of water surface covered by at least 0.01 g/m ²	High
4B-2: Water Surface Degree SRAR Oiling	Low	Medium	High	The mean area of water contaminated above 0.01 g/m ² was 597,000 mi ²	
4C-1: Shoreline Probability SRAR Oiling	Low	Medium	High	12% of the model runs resulted in shoreline oiling of 1 g/m ²	High
4C-2: Shoreline Degree SRAR Oiling	Low	Medium	High	The length of shoreline contaminated by at least 1 g/m ² was 139 mi	

For the Most Probable Discharge of 13,200 bbl, the socio-economic risk from potential releases of crude oil from the *Esso Gettysburg* is summarized as listed below and indicated in the far-right column in Table 4-3:

- Water column resources – Medium, because a moderate area of water column would be impacted in areas with fishing grounds and other offshore resources (e.g., marine sanctuary)
- Water surface resources – High, because a large area of water surface would be impacted offshore in areas with shipping lanes and other offshore activities. It should be noted that oil on the surface will not be continuous but rather be broken and patchy and in the form of sheens, tarballs, and streamers
- Shoreline resources – High, because a relatively large length of shoreline would be impacted in areas with high-value and sensitive resources

Table 4-3: Socio-economic risk factor ranks for the **Most Probable Discharge of 13,200 bbl** of crude oil from the *Esso Gettysburg*.

Risk Factor	Risk Score			Explanation of Risk Score	Final Score
4A-1: Water Column Probability SRAR Oiling	Low	Medium	High	100% of the model runs resulted in at least 0.2 mi ² of the upper 33 feet of the water column contaminated above 1 ppb aromatics	Med
4A-2: Water Column Degree SRAR Oiling	Low	Medium	High	The mean volume of water contaminated above 1 ppb was 81 mi ² of the upper 33 feet of the water column	
4B-1: Water Surface Probability SRAR Oiling	Low	Medium	High	100% of the model runs resulted in at least 1,000 mi ² of water surface covered by at least 0.01 g/m ²	High
4B-2: Water Surface Degree SRAR Oiling	Low	Medium	High	The mean area of water contaminated above 0.01 g/m ² was 152,400 mi ²	
4C-1: Shoreline Probability SRAR Oiling	Low	Medium	High	14% of the model runs resulted in shoreline oiling of 1 g/m ²	High
4C-2: Shoreline Degree SRAR Oiling	Low	Medium	High	The length of shoreline contaminated by at least 1 g/m ² was 104 mi	

SECTION 5: OVERALL RISK ASSESSMENT AND RECOMMENDATIONS FOR ASSESSMENT, MONITORING, OR REMEDIATION

The overall risk assessment for the *Esso Gettysburg* is comprised of a compilation of several components that reflect the best available knowledge about this particular site. Those components are reflected in the previous sections of this document and are:

- Vessel casualty information and how the site formation processes have worked on this vessel
- Ecological resources at risk
- Socio-economic resources at risk
- Other complicating factors (war graves, other hazardous cargo, etc.)

Table 5-1 summarizes the screening-level risk assessment scores for the different risk factors, as discussed in the previous sections. The ecological and socio-economic risk factors are presented as a single score for water column, water surface, and shoreline resources as the scores were consolidated for each element. For the ecological and socio-economic risk factors each has two components, probability and degree. Of those two, degree is given more weight in deciding the combined score for an individual factor, e.g., a high probability and medium degree score would result in a medium overall for that factor.

In order to make the scoring more uniform and replicable between wrecks, a value was assigned to each of the 7 criteria. This assessment has a total of 7 criteria (based on table 5-1) with 3 possible scores for each criteria (L, M, H). Each was assigned a point value of L=1, M=2, H=3. The total possible score is 21 points, and the minimum score is 7. The resulting category summaries are:

Low Priority	7-11
Medium Priority	12-14
High Priority	15-21

For the Worst Case Discharge, *Esso Gettysburg* scores High with 18 points; for the Most Probable Discharge, *Esso Gettysburg* also scores High with 16 points. Under the National Contingency Plan, the U.S. Coast Guard and the Regional Response Team have the primary authority and responsibility to plan, prepare for, and respond to oil spills in U.S. waters. Based on the technical review of available information, NOAA proposes the following recommendations for the *Esso Gettysburg*. The final determination of what type of action, if any, rests with the U.S. Coast Guard.

<i>Esso Gettysburg</i>	Possible NOAA Recommendations
	Wreck should be considered for further assessment to determine the vessel condition, amount of oil onboard, and feasibility of oil removal action
✓	Location is unknown; Use surveys of opportunity to attempt to locate this vessel and gather more information on the vessel condition
	Conduct active monitoring to look for releases or changes in rates of releases
✓	Be noted in the Area Contingency Plans so that if a mystery spill is reported in the general area, this vessel could be investigated as a source
✓	Conduct outreach efforts with the technical and recreational dive community as well as commercial and recreational fishermen who frequent the area, to gain awareness of changes in the site

Table 5-1: Summary of risk factors for the *Esso Gettysburg*.

Vessel Risk Factors		Data Quality Score	Comments	Risk Score	
Pollution Potential Factors	A1: Oil Volume (total bbl)	Medium	Maximum of 131,146 bbl, no known leakage	Med	
	A2: Oil Type	Medium	Cargo is West Texas crude oil, a Group III oil type		
	B: Wreck Clearance	High	No		
	C1: Burning of the Ship	High	Yes		
	C2: Oil on Water	High	Yes		
	D1: Nature of Casualty	High	Two torpedoes, explosion and fire		
	D2: Structural Breakup	High	Unknown		
Archaeological Assessment	Archaeological Assessment	High	Detailed sinking reports exist, assessment is believed to be very accurate	Not Scored	
Operational Factors	Wreck Orientation	Low	Unknown	Not Scored	
	Depth	Low	Unknown		
	Visual or Remote Sensing Confirmation of Site Condition	Low	Unknown		
	Other Hazardous Materials Onboard	High	No		
	Munitions Onboard	High	one 4in gun, one 3in gun, eight 20mm guns		
	Gravesite (Civilian/Military)	High	Yes		
	Historical Protection Eligibility (NHPA/SMCA)	High	NHPA and possibly SMCA		
				WCD	Most Probable
Ecological Resources	3A: Water Column Resources	High	Areas above thresholds are in open water settings where most resources are highly mobile and able to avoid exposure	Med	Med
	3B: Water Surface Resources	High	Crude oils weather to persistent tarballs that can travel long distances posing risks to the many birds and sea turtles in the area, esp. when concentrated in convergence zones and <i>Sargassum</i>	High	High
	3C: Shore Resources	High	Persistent tarballs strand mostly on beaches but also marshes and tidal flats, fouling habitats and animals	Med	Low
Socio-Economic Resources	4A: Water Column Resources	High	Large to moderate area of water column would be impacted in important fishing grounds and marine sanctuary	High	Med
	4B: Water Surface Resources	High	A large area of water surface would be impacted offshore in areas with shipping lanes and other offshore activities	High	High
	4C: Shore Resources	High	A relatively large length of shoreline would be impacted in areas with high-value and sensitive resources	High	High
Summary Risk Scores				18	16