



Screening Level Risk Assessment Package

Taborfjell



ENVIRONMENTAL
RESEARCH
CONSULTING

National Oceanic and
Atmospheric Administration

Office of National Marine Sanctuaries
Daniel J. Basta, Director
Lisa Symons
John Wagner

Office of Response and Restoration
Dave Westerholm, Director
Debbie Payton
Doug Helton



Table of Contents

Project Background	ii
Executive Summary	1
Section 1: Vessel Background Information: Remediation of Underwater Legacy	
Environmental Threats (RULET).....	2
Vessel Particulars.....	2
Casualty Information.....	3
Wreck Location.....	4
Casualty Narrative	4
General Notes	4
Wreck Condition/Salvage History	5
Archaeological Assessment	5
Assessment.....	5
Background Information References	6
Vessel Risk Factors.....	6
Section 2: Environmental Impact Modeling.....	13
Release Scenarios Used in the Modeling	13
Oil Type for Release	15
Oil Thickness Thresholds.....	15
Potential Impacts to the Water Column.....	16
Potential Water Surface Slick.....	17
Potential Shoreline Impacts.....	19
Section 3: Ecological Resources At Risk	20
Ecological Risk Factors	23
Section 4: Socio-Economic Resources At Risk	28
Socio-Economic Risk Factors	31
Section 5: Overall Risk Assessment and Recommendations for Assessment,	
Monitoring, or Remediation	36

Project Background

The past century of commerce and warfare has left a legacy of thousands of sunken vessels along the U.S. coast. Many of these wrecks pose environmental threats because of the hazardous nature of their cargoes, presence of munitions, or bunker fuel oils left onboard. As these wrecks corrode and decay, they may release oil or hazardous materials. Although a few vessels, such as USS *Arizona* in Hawaii, are well-publicized environmental threats, most wrecks, unless they pose an immediate pollution threat or impede navigation, are left alone and are largely forgotten until they begin to leak.

In order to narrow down the potential sites for inclusion into regional and area contingency plans, in 2010, Congress appropriated \$1 million to identify the most ecologically and economically significant potentially polluting wrecks in U.S. waters. This project supports the U.S. Coast Guard and the Regional Response Teams as well as NOAA in prioritizing threats to coastal resources while at the same time assessing the historical and cultural significance of these nonrenewable cultural resources.


The potential polluting shipwrecks were identified through searching a broad variety of historical sources. NOAA then worked with Research Planning, Inc., RPS ASA, and Environmental Research Consulting to conduct the modeling forecasts, and the ecological and environmental resources at risk assessments.

Initial evaluations of shipwrecks located within American waters found that approximately 600-1,000 wrecks could pose a substantial pollution threat based on their age, type and size. This includes vessels sunk after 1891 (when vessels began being converted to use oil as fuel), vessels built of steel or other durable material (wooden vessels have likely deteriorated), cargo vessels over 1,000 gross tons (smaller vessels would have limited cargo or bunker capacity), and any tank vessel.

Additional ongoing research has revealed that 87 wrecks pose a potential pollution threat due to the violent nature in which some ships sank and the structural reduction and demolition of those that were navigational hazards. To further screen and prioritize these vessels, risk factors and scores have been applied to elements such as the amount of oil that could be on board and the potential ecological or environmental impact.

Executive Summary: *Taborfjell*

The freighter *Taborfjell*, torpedoed and sunk during World War II off the coast of Massachusetts in 1942, was identified as a potential pollution threat, thus a screening-level risk assessment was conducted. The different sections of this document summarize what is known about the *Taborfjell*, the results of environmental impact modeling composed of different release scenarios, the ecological and socio-economic resources that would be at risk in the event of releases, the screening-level risk scoring results and overall risk assessment, and recommendations for assessment, monitoring, or remediation.



Based on this screening-level assessment, each vessel was assigned a summary score calculated using the seven risk criteria described in this report. For the Worst Case Discharge, *Taborfjell* scores Low with 11 points; for the Most Probable Discharge (10% of the Worst Case volume), *Taborfjell* also scores Low with 11 points. Given these scores, the medium/low level of data certainty, and that the location of this vessel is unknown, NOAA recommends that general notations are made in the Area Contingency Plans so that if a mystery spill is reported in the general area, this vessel could be investigated as a source. Outreach efforts with the technical and recreational dive community as well as commercial and recreational fishermen who frequent the area would be helpful to gain awareness of localized spills in the general area where the vessel is believed lost.

Vessel Risk Factors		Risk Score	
Pollution Potential Factors	A1: Oil Volume (total bbl)	Med	
	A2: Oil Type		
	B: Wreck Clearance		
	C1: Burning of the Ship		
	C2: Oil on Water		
	D1: Nature of Casualty		
Archaeological Assessment	D2: Structural Breakup	Not Scored	
	Archaeological Assessment		
Operational Factors	Wreck Orientation	Not Scored	
	Depth		
	Confirmation of Site Condition		
	Other Hazardous Materials		
	Munitions Onboard		
	Gravesite (Civilian/Military)		
Historical Protection Eligibility			
		WCD	MP (10%)
Ecological Resources	3A: Water Column Resources	Low	Low
	3B: Water Surface Resources	Med	Med
	3C: Shore Resources	Med	Low
Socio-Economic Resources	4A: Water Column Resources	Low	Low
	4B: Water Surface Resources	Low	Low
	4C: Shore Resources	Med	Med
Summary Risk Scores		11	11

The determination of each risk factor is explained in the document. This summary table is found on page 37.

SECTION 1: VESSEL BACKGROUND INFORMATION: REMEDIATION OF UNDERWATER LEGACY ENVIRONMENTAL THREATS (RULET)

Vessel Particulars

Official Name: *Taborfjell*

Official Number: Unknown

Vessel Type: Freighter

Vessel Class: Unknown

Former Names: N/A

Year Built: 1938

Builder: Nylands Verksted, Oslo

Builder's Hull Number: Unknown

Flag: Norwegian

Owner at Loss: Norwegian Shipping and Trade Mission

Controlled by: Unknown

Chartered to: Unknown

Operated by: Unknown

Homeport: Oslo

Length: 250 feet

Beam: 41 feet

Depth: 14 feet

Gross Tonnage: 1,339

Net Tonnage: 742

Hull Material: Steel

Hull Fastenings: Riveted

Powered by: Oil-fired steam

Bunker Type: Heavy fuel oil (Bunker C)

Bunker Capacity (bbl): Unknown

Average Bunker Consumption (bbl) per 24 hours: Unknown

Liquid Cargo Capacity (bbl): Unknown

Dry Cargo Capacity: Unknown

Tank or Hold Description: Unknown

Casualty Information

Port Departed: New York

Destination Port: Montreal, Canada

Date Departed: April 28, 1942

Date Lost: April 30, 1942

Number of Days Sailing: ≈ 3

Cause of Sinking: Act of War (Torpedoes)

Latitude (DD): 42.3334

Longitude (DD): -69.1661

Nautical Miles to Shore: 54

Nautical Miles to NMS: 54

Nautical Miles to MPA: 0

Nautical Miles to Fisheries: Unknown

Approximate Water Depth (Ft): 800

Bottom Type: Unknown

Is There a Wreck at This Location? Unknown, the wreck has never been located or surveyed

Wreck Orientation: Unknown

Vessel Armament: One 3-in gun, one Colt .50-mm gun, two Hotchkiss guns, two Lewis machine guns, and one Marlin machine gun

Cargo Carried when Lost: 2,200 tons of unrefined sugar in 16,200 bags, all holds filled to the hatch coamings

Cargo Oil Carried (bbl): 0

Cargo Oil Type: N/A

Probable Fuel Oil Remaining (bbl): Unknown, $\leq 8,000$

Fuel Type: Heavy fuel oil (Bunker C)

Total Oil Carried (bbl): $\leq 8,000$ (likely far less)

Dangerous Cargo or Munitions: Yes

Munitions Carried: Munitions for onboard weapons

Demolished after Sinking: No

Salvaged: No

Cargo Lost: Yes

Reportedly Leaking: No

Historically Significant: Yes

Gravesite: Yes

Salvage Owner: Not known if any

Wreck Location




Chart Number: 13003

Casualty Narrative

"At 07.37 hours on 30 Apr, 1942, the unescorted *Taborfjell* (Master Erling Nilsen) was hit by two torpedoes from *U-576* about 95 miles east of Cape Cod and caused the ship to sink within one minute. The ship sank so quickly that no lifeboats could be launched and the three survivors had all jumped overboard and rescued themselves on a raft. They observed how the U-boat passed them without taking any action and had seen three other survivors but they later disappeared. After 20 hours, the survivors were picked up by HMS *P-552* (P 552) and landed at St. Johns on 7 May."

<http://www.uboot.net:8080/allies/merchants/ships/1575.html>

General Notes

AWOIS Data:

DESCRIPTION

24 NO.201; CARGO, 1339 GT, SUNK 4/30/42 BY SUBMARINE, POSITION ACCURACY 1-3 MILES; REPORTED IN 135 FATHOMS THRU ROYAL CANADIAN NAVY-WK LIST OF 12/9/42.

27 NO.724; FTR, 1339 NT, IN 135 FMS OF WATER, REPORTED THRU THE ROYAL CANADIAN NAVY WK LIST OF 12/9/42.

SURVEY REQUIREMENTS MISCELLANEOUS; NONE, EXCESSIVE DEPTH.

Wreck Condition/Salvage History

Unknown; the wreck has never been located or surveyed.

Archaeological Assessment

The archaeological assessment provides additional primary source based documentation about the sinking of vessels. It also provides condition-based archaeological assessment of the wrecks when possible. It does not provide a risk-based score or definitively assess the pollution risk or lack thereof from these vessels, but includes additional information that could not be condensed into database form.

Where the current condition of a shipwreck is not known, data from other archaeological studies of similar types of shipwrecks provide the means for brief explanations of what the shipwreck might look like and specifically whether it is thought there is sufficient structural integrity to retain oil. This is more subjective than the Pollution Potential Tree and computer-generated resource at risk models, and as such provides an additional viewpoint to examine risk assessments and assess the threat posed by these shipwrecks. It also addresses questions of historical significance and the relevant historic preservation laws and regulations that will govern on-site assessments.

In some cases where little additional historic information has been uncovered about the loss of a vessel, archaeological assessments cannot be made with any degree of certainty and were not prepared. For vessels with full archaeological assessments, NOAA archaeologists and contracted archivists have taken photographs of primary source documents from the National Archives that can be made available for future research or on-site activities.

Assessment

The wreck of *Taborfjell* has never been located so there are no site reports that would allow NOAA archaeologists to provide a condition based archaeological assessment of the shipwreck. Some additional analysis can be made based on the historic sinking reports of the ship that may be of utility to the U.S. Coast Guard. We do know from archival research that the ship was struck by one torpedo (possibly two) on the starboard side. The torpedo struck between the No. 3 and No. 4 hatches and ripped open a hole on the starboard side of the ship to the stern, leaving a gaping hole above the waterline about 40 to 50 feet long and about 10 feet deep, extending up to the deck plates. This extensive damage caused the ship to sink by the stern within two minutes.

Based on the location of the torpedo damage, the vessel's bunker tanks may not have been damaged by the initial blast since the engine room on this ship is believed to have been located amidships. This ship is one of the smallest ships in the RULET database, and it is likely that the ship had a relatively small bunker capacity. Because the shipwreck has never been discovered, it is not possible to determine with any degree of accuracy what the current condition of the wreck is and how likely the vessel is to contain oil.

It should be noted, however, that there is a great amount of discrepancy between the reported sinking locations for this ship. The coordinates used to map this wreck place the wreck approximately 54 miles offshore, while some of the historic sinking reports state that the vessel was lost approximately 100 miles

due east of Cape Cod. Based on the large degree of inaccuracy between these sinking locations, it is unlikely that the shipwreck will be intentionally located.

Should the vessel be located in a survey of opportunity or due to a mystery spill attributed to this vessel, it should be noted that this vessel is of historic significance and will require appropriate actions be taken under the National Historic Preservation Act (NHPA) and the Sunken Military Craft Act (SMCA) prior to any actions that could impact the integrity of the vessel. This vessel may be eligible for listing on the National Register of Historic Places. The site is also considered a war grave and appropriate actions should be undertaken to minimize disturbance to the site.

Background Information References

Vessel Image Sources: No image of this vessel has been located by NOAA to date

Construction Diagrams or Plans in RULET Database? No

Text References:

-Office of the Chief of Naval Operations

Tenth Fleet ASW Analysis & Stat. Section Series XIII. Report and Analyses of U. S. and Allied Merchant Shipping Losses 1941-1945 Stocco – Thirlby, Records of the Office of the Chief of Naval Operations, Box 249, Record Group 38, National Archives at College Park, MD.

-AWOIS database

-<http://www.uboaat.net/allies/merchants/ships/1575.html>

Vessel Risk Factors

In this section, the risk factors that are associated with the vessel are defined and then applied to the *Taborfjell* based on the information available. These factors are reflected in the pollution potential risk assessment development by the U.S. Coast Guard Salvage Engineering Response Team (SERT) as a means to apply a salvage engineer's perspective to the historical information gathered by NOAA. This analysis reflected in Figure 1-1 is simple and straightforward and, in combination with the accompanying archaeological assessment, provides a picture of the wreck that is as complete as possible based on current knowledge and best professional judgment. This assessment *does not* take into consideration operational constraints such as depth or unknown location, but rather attempts to provide a replicable and objective screening of the historical date for each vessel. SERT reviewed the general historical information available for the database as a whole and provided a stepwise analysis for an initial indication of Low/Medium/High values for each vessel.

In some instances, nuances from the archaeological assessment may provide additional input that will amend the score for Section 1. Where available, additional information that may have bearing on operational considerations for any assessment or remediation activities is provided.




Figure 1-1: U.S. Coast Guard Salvage Engineering Response Team (SERT) developed the above Pollution Potential Decision Tree.

Each risk factor is characterized as High, Medium, or Low Risk or a category-appropriate equivalent such as No, Unknown, Yes, or Yes Partially. The risk categories correlate to the decision points reflected in Figure 1-1.

Each of the risk factors also has a “data quality modifier” that reflects the completeness and reliability of the information on which the risk ranks were assigned. The quality of the information is evaluated with

respect to the factors required for a reasonable preliminary risk assessment. The data quality modifier scale is:

- **High Data Quality:** All or most pertinent information on wreck available to allow for thorough risk assessment and evaluation. The data quality is high and confirmed.
- **Medium Data Quality:** Much information on wreck available, but some key factor data are missing or the data quality is questionable or not verified. Some additional research needed.
- **Low Data Quality:** Significant issues exist with missing data on wreck that precludes making preliminary risk assessment, and/or the data quality is suspect. Significant additional research needed.

In the following sections, the definition of low, medium, and high for each risk factor is provided. Also, the classification for the *Taborfjell* is provided, both as text and as shading of the applicable degree of risk bullet.

Pollution Potential Factors

Risk Factor A1: Total Oil Volume

The oil volume classifications correspond to the U.S. Coast Guard spill classifications:

- **Low Volume: Minor Spill** <240 bbl (10,000 gallons)
- **Medium Volume: Medium Spill** ≥240 – 2,400 bbl (100,000 gallons)
- **High Volume: Major Spill** ≥2,400 bbl (≥100,000 gallons)

The oil volume risk classifications refer to the volume of the most-likely Worst Case Discharge from the vessel and are based on the amount of oil believed or confirmed to be on the vessel.

The *Taborfjell* is ranked as High Volume because it is thought to have a potential for up to 3,000 bbl (based on known fuel capacities for vessels of similar size), although some of that may have been lost at the time of the casualty due to the explosion. Data quality is low.

The risk factor for volume also incorporates any reports or anecdotal evidence of actual leakage from the vessel or reports from divers of oil in the overheads, as opposed to potential leakage. This reflects the history of the vessel's leakage. There are no reports of leakage from the *Taborfjell*.

Risk Factor A2: Oil Type

The oil type(s) on board the wreck are classified only with regard to persistence, using the U.S. Coast Guard oil grouping¹. (Toxicity is dealt with in the impact risk for the Resources at Risk classifications.) The three oil classifications are:

- **Low Risk: Group I Oils** – non-persistent oil (e.g., gasoline)

¹ Group I Oil or Nonpersistent oil is defined as "a petroleum-based oil that, at the time of shipment, consists of hydrocarbon fractions: At least 50% of which, by volume, distill at a temperature of 340°C (645°F); and at least 95% of which, by volume, distill at a temperature of 370°C (700°F)."

Group II - Specific gravity less than 0.85 crude [API° >35.0]

Group III - Specific gravity between 0.85 and less than .95 [API° ≤35.0 and >17.5]

Group IV - Specific gravity between 0.95 to and including 1.0 [API° ≤17.5 and >10.0]

- **Medium Risk: Group II – III Oils** – medium persistent oil (e.g., diesel, No. 2 fuel, light crude, medium crude)
- **High Risk: Group IV** – high persistent oil (e.g., heavy crude oil, No. 6 fuel oil, Bunker C)

The *Taborfjell* is classified as High Risk because the bunker oil is heavy fuel oil, a Group IV oil type. Data quality is high.

Was the wreck demolished?

Risk Factor B: Wreck Clearance

This risk factor addresses whether or not the vessel was historically reported to have been demolished as a hazard to navigation or by other means such as depth charges or aerial bombs. This risk factor is based on historic records and does not take into account what a wreck site currently looks like. The risk categories are defined as:

- **Low Risk:** The wreck was reported to have been entirely destroyed after the casualty
- **Medium Risk:** The wreck was reported to have been partially cleared or demolished after the casualty
- **High Risk:** The wreck was not reported to have been cleared or demolished after the casualty
- **Unknown:** It is not known whether or not the wreck was cleared or demolished at the time of or after the casualty

The *Taborfjell* is classified as High Risk because there are no known historic accounts of the wreck being demolished as a hazard to navigation. Data quality is high.

Was significant cargo or bunker lost during casualty?

Risk Factor C1: Burning of the Ship

This risk factor addresses any burning that is known to have occurred at the time of the vessel casualty and may have resulted in oil products being consumed or breaks in the hull or tanks that would have increased the potential for oil to escape from the shipwreck. The risk categories are:

- **Low Risk:** Burned for multiple days
- **Medium Risk:** Burned for several hours
- **High Risk:** No burning reported at the time of the vessel casualty
- **Unknown:** It is not known whether or not the vessel burned at the time of the casualty

The *Taborfjell* is classified as High Risk because there was no report of fire at the time of casualty. Data quality is high.

Risk Factor C2: Reported Oil on the Water

This risk factor addresses reports of oil on the water at the time of the vessel casualty. The amount is relative and based on the number of available reports of the casualty. Seldom are the reports from trained observers so this is very subjective information. The risk categories are defined as:

- **Low Risk:** Large amounts of oil reported on the water by multiple sources
- **Medium Risk:** Moderate to little oil reported on the water during or after the sinking event

- **High Risk:** No oil reported on the water
- **Unknown:** It is not known whether or not there was oil on the water at the time of the casualty

The *Taborfjell* is classified as High Risk because no oil is known to have been reported spreading across the water as the vessel went down. Data quality is high.

Is the cargo area damaged?

Risk Factor D1: Nature of the Casualty

This risk factor addresses the means by which the vessel sank. The risk associated with each type of casualty is determined by the how violent the sinking event was and the factors that would contribute to increased initial damage or destruction of the vessel (which would lower the risk of oil, other cargo, or munitions remaining on board). The risk categories are:

- **Low Risk:** Multiple torpedo detonations, multiple mines, severe explosion
- **Medium Risk:** Single torpedo, shellfire, single mine, rupture of hull, breaking in half, grounding on rocky shoreline
- **High Risk:** Foul weather, grounding on soft bottom, collision
- **Unknown:** The cause of the loss of the vessel is not known

The *Taborfjell* is classified as Medium Risk because the survivor reports state there was one torpedo blast, but the article available on uboat.net states that there were two torpedo detonations. Data quality is medium.

Risk Factor D2: Structural Breakup

This risk factor takes into account how many pieces the vessel broke into during the sinking event or since sinking. This factor addresses how likely it is that multiple components of a ship were broken apart including tanks, valves, and pipes. Experience has shown that even vessels broken in three large sections can still have significant pollutants on board if the sections still have some structural integrity. The risk categories are:

- **Low Risk:** The vessel is broken into more than three pieces
- **Medium Risk:** The vessel is broken into two-three pieces
- **High Risk:** The vessel is not broken and remains as one contiguous piece
- **Unknown:** It is currently not known whether or not the vessel broke apart at the time of loss or after sinking

The *Taborfjell* is classified as Unknown Risk because it is not known if the vessel broken apart at the time of casualty or after it sank since the location is unknown. Data quality is low.

Factors That May Impact Potential Operations

Orientation (degrees)

This factor addresses what may be known about the current orientation of the intact pieces of the wreck (with emphasis on those pieces where tanks are located) on the seafloor. For example, if the vessel turtled,

not only may it have avoided demolition as a hazard to navigation, but it has a higher likelihood of retaining an oil cargo in the non-vented and more structurally robust bottom of the hull.

The location of the *Taborfjell* is unknown. Data quality is low.

Depth

Depth information is provided where known. In many instances, depth will be an approximation based on charted depths at the last known locations.

The depth for *Taborfjell* is believed to be greater than 800 feet. Data quality is low.

Visual or Remote Sensing Confirmation of Site Condition

This factor takes into account what the physical status of wreck site as confirmed by remote sensing or other means such as ROV or diver observations and assesses its capability to retain a liquid cargo. This assesses whether or not the vessel was confirmed as entirely demolished as a hazard to navigation, or severely compromised by other means such as depth charges, aerial bombs, or structural collapse.

The location of the *Taborfjell* is unknown. Data quality is low.

Other Hazardous (Non-Oil) Cargo on Board

This factor addresses hazardous cargo other than oil that may be on board the vessel and could potentially be released, causing impacts to ecological and socio-economic resources at risk.

There are no reports of hazardous materials onboard. Data quality is high.

Munitions on Board

This factor addresses hazardous cargo other than oil that may be on board the vessel and could potentially be released or detonated causing impacts to ecological and socio-economic resources at risk.

The *Taborfjell* had munitions for onboard weapons, one 3-in gun, one Colt .50-mm gun, two Hotchkiss guns, two Lewis machine guns, and one Marlin machine gun. Data quality is high.

Vessel Pollution Potential Summary

Table 1-1 summarizes the risk factor scores for the pollution potential and mitigating factors that would reduce the pollution potential for the *Taborfjell*. Operational factors are listed but do not have a risk score.

Table 1-1: Summary matrix for the vessel risk factors for the *Taborfjell* color-coded as red (high risk), yellow (medium risk), and green (low risk).

Vessel Risk Factors		Data Quality Score	Comments	Risk Score
Pollution Potential Factors	A1: Oil Volume (total bbl)	Low	Maximum of 3,000 bbl, not reported to be leaking	Med
	A2: Oil Type	High	Bunker oil is heavy fuel oil, a Group IV oil type	
	B: Wreck Clearance	High	Vessel not reported as cleared	
	C1: Burning of the Ship	High	No fire was reported	
	C2: Oil on Water	High	No oil was reported on the water	
	D1: Nature of Casualty	Medium	One (possibly two torpedo detonations)	
	D2: Structural Breakup	Low	Unknown structural breakup	
Archaeological Assessment	Archaeological Assessment	Low	Limited sinking records of this ship exist, and the site has never been located, assessment is believed to have low accuracy	Not Scored
Operational Factors	Wreck Orientation	Low	Unknown, potential to be upright	Not Scored
	Depth	Low	> 800 ft	
	Visual or Remote Sensing Confirmation of Site Condition	Low	Location unknown	
	Other Hazardous Materials Onboard	High	No	
	Munitions Onboard	High	Munitions for onboard weapons	
	Gravesite (Civilian/Military)	High	Yes	
	Historical Protection Eligibility (NHPA/SMCA)	High	NHPA and possibly SMCA	

SECTION 2: ENVIRONMENTAL IMPACT MODELING

To help evaluate the potential transport and fates of releases from sunken wrecks, NOAA worked with RPS ASA to run a series of generalized computer model simulations of potential oil releases. The results are used to assess potential impacts to ecological and socio-economic resources, as described in Sections 3 and 4. The modeling results are useful for this screening-level risk assessment; however, it should be noted that detailed site/vessel/and seasonally specific modeling would need to be conducted prior to any intervention on a specific wreck.

Release Scenarios Used in the Modeling

The potential volume of leakage at any point in time will tend to follow a probability distribution. Most discharges are likely to be relatively small, though there could be multiple such discharges. There is a lower probability of larger discharges, though these scenarios would cause the greatest damage. A **Worst Case Discharge** (WCD) would involve the release of all of the cargo oil and bunkers present on the vessel. In the case of the *Taborfjell* this would be about 3,000 bbl of Bunker C fuel oil based on current estimates of the maximum amount of oil remaining onboard the wreck.

The likeliest scenario of oil release from most sunken wrecks, including the *Taborfjell*, is a small, episodic release that may be precipitated by disturbance of the vessel in storms. Each of these episodic releases may cause impacts and require a response. **Episodic** releases are modeled using 1% of the WCD. Another scenario is a very low chronic release, i.e., a relatively regular release of small amounts of oil that causes continuous oiling and impacts over the course of a long period of time. This type of release would likely be precipitated by corrosion of piping that allows oil to flow or bubble out at a slow, steady rate. **Chronic** releases are modeled using 0.1% of the WCD.

The **Most Probable** scenario is premised on the release of all the oil from one tank. In the absence of information on the number and condition of the cargo or fuel tanks for all the wrecks being assessed, this scenario is modeled using 10% of the WCD. The **Large** scenario is loss of 50% of the WCD. The five major types of releases are summarized in Table 2-1. The actual type of release that occurs will depend on the condition of the vessel, time factors, and disturbances to the wreck. Note that episodic and chronic release scenarios represent a small release that is repeated many times, potentially repeating the same magnitude and type of impact(s) with each release. The actual impacts would depend on the environmental factors such as real-time and forecast winds and currents during each release and the types/quantities of ecological and socio-economic resources present.

The model results here are based on running the RPS ASA Spill Impact Model Application Package (SIMAP) two hundred times for each of the five spill volumes shown in Table 2-1. The model randomly selects the date of the release, and corresponding environmental, wind, and ocean current information from a long-term wind and current database.

When a spill occurs, the trajectory, fate, and effects of the oil will depend on environmental variables, such as the wind and current directions over the course of the oil release, as well as seasonal effects. The

magnitude and nature of potential impacts to resources will also generally have a strong seasonal component (e.g., timing of bird migrations, turtle nesting periods, fishing seasons, and tourism seasons).

Table 2-1: Potential oil release scenario types for the *Taborfjell*.

Scenario Type	Release per Episode	Time Period	Release Rate	Relative Likelihood	Response Tier
Chronic (0.1% of WCD)	3 bbl	Fairly regular intervals or constant	100 bbl over several days	More likely	Tier 1
Episodic (1% of WCD)	30 bbl	Irregular intervals	Over several hours or days	Most Probable	Tier 1-2
Most Probable (10% of WCD)	300 bbl	One-time release	Over several hours or days	Most Probable	Tier 2
Large (50% of WCD)	1,500 bbl	One-time release	Over several hours or days	Less likely	Tier 2-3
Worst Case	3,000 bbl	One-time release	Over several hours or days	Least likely	Tier 3

The modeling results represent 200 simulations for each spill volume with variations in spill trajectory based on winds and currents. The spectrum of the simulations gives a perspective on the variations in likely impact scenarios. Some resources will be impacted in nearly all cases; some resources may not be impacted unless the spill trajectory happens to go in that direction based on winds and currents at the time of the release and in its aftermath.

For the large and WCD scenarios, the duration of the release was assumed to be 12 hours, envisioning a storm scenario where the wreck is damaged or broken up, and the model simulations were run for a period of 30 days. The releases were assumed to be from a depth between 2-3 meters above the sea floor, using the information known about the wreck location and depth.

As discussed in the NOAA 2013 Risk Assessment for Potentially Polluting Wrecks in U.S. Waters, NOAA identified 70 high and medium priority wrecks for screening-level risk assessment. Within the available funds, it was not feasible to conduct computer model simulations of all 70 high and medium priority wrecks. Therefore, efforts were made to create “clusters” of vessels in reasonable proximity and with similar oil types. In general, the wreck with the largest potential amount of oil onboard was selected for modeling of oil release volumes, and the results were used as surrogates for the other vessels in the cluster. In particular, the regression curves created for the modeled wreck were used to determine the impacts to water column, water surface, and shoreline resources. The *Taborfjell*, with up to 3,000 bbl of heavy fuel onboard, was clustered with the *Cherokee*, which was modeled at 10,000 bbl of heavy fuel oil. Figure 2-1 shows the location of both vessels.

It is important to acknowledge that these scenarios are only for this screening-level assessment. Detailed site/vessel/and seasonally specific modeling would need to be conducted prior to any intervention on a specific wreck.




Figure 2-1: Location of the *Taborfjell* (red triangle), the wreck discussed in this package, and the *Cherokee* (red circle) which was the wreck that was actually modeled in the computer modeling simulations. The results for the *Cherokee* are used to estimate the impacts of releases from the *Taborfjell*, as discussed in the text.

Oil Type for Release

The *Taborfjell* contained a maximum of 3,000 bbl of heavy fuel oil (a Group IV oil). Thus, the spill model for the *Cherokee*, which was run using heavy fuel oil, was used for this scoping assessment of the *Taborfjell*.

Oil Thickness Thresholds

The model results are reported for different oil thickness thresholds, based on the amount of oil on the water surface or shoreline and the resources potentially at risk. Table 2-2 shows the terminology and thicknesses used in this report, for both oil thickness on water and the shoreline. For oil on the water surface, a thickness of 0.01 g/m^2 , which would appear as a barely visible sheen, was used as the threshold for socio-economic impacts because often fishing is prohibited in areas with any visible oil, to prevent contamination of fishing gear and catch. A thickness of 10 g/m^2 was used as the threshold for ecological impacts, primarily due to impacts to birds, because that amount of oil has been observed to be enough to mortally impact birds and other wildlife. In reality, it is very unlikely that oil would be evenly distributed on the water surface. Spilled oil is always distributed patchily on the water surface in bands or tarballs with clean water in between. So, Table 2-2a shows the number of tarballs per acre on the water surface for these oil thickness thresholds, assuming that each tarball was a sphere that was 1 inch in diameter. For oil stranded onshore, a thickness of 1 g/m^2 was used as the threshold for socio-economic impacts because that amount of oil would conservatively trigger the need for shoreline cleanup on amenity beaches. A thickness of 100 g/m^2 was used as the threshold for ecological impacts based on a synthesis of

the literature showing that shoreline life has been affected by this degree of oiling.² Because oil often strands onshore as tarballs, Table 2-2a shows the number of tarballs per m² on the shoreline for these oil thickness thresholds, assuming that each tarball was a sphere that was 1 inch in diameter.

Table 2-2a: Oil thickness thresholds used in calculating area of water impacted. Refer to Sections 3 and 4 for explanations of the thresholds for ecological and socio-economic resource impacts.

Oil Description	Sheen Appearance	Approximate Sheen Thickness		No. of 1 inch Tarballs	Threshold/Risk Factor
Oil Sheen	Barely Visible	0.00001 mm	0.01 g/m ²	~5-6 tarballs per acre	Socio-economic Impacts to Water Surface/Risk Factor 4B-1 and 2
Heavy Oil Sheen	Dark Colors	0.01 mm	10 g/m ²	~5,000-6,000 tarballs per acre	Ecological Impacts to Water Surface/ Risk Factor 3B-1 and 2

Table 2-2b: Oil thickness thresholds used in calculating miles of shoreline impacted. Refer to Sections 3 and 4 for explanations of the thresholds for ecological and socio-economic resource impacts.

Oil Description	Oil Appearance	Approximate Sheen Thickness		No. of 1 inch Tarballs	Threshold/Risk Factor
Oil Sheen/Tarballs	Dull Colors	0.001 mm	1 g/m ²	~0.12-0.14 tarballs/m ²	Socio-economic Impacts to Shoreline Users/Risk Factor 4C-1 and 2
Oil Slick/Tarballs	Brown to Black	0.1 mm	100 g/m ²	~12-14 tarballs/m ²	Ecological Impacts to Shoreline Habitats/Risk Factor 3C-1 and 2

Potential Impacts to the Water Column

Impacts to the water column from an oil release from the *Taborfjell* will be determined by the volume of leakage. Because oil from sunken vessels will be released at low pressures, the droplet sizes will be large enough for the oil to float to the surface. Therefore, impacts to water column resources will result from the natural dispersion of the floating oil slicks on the surface, which is limited to about the top 33 feet. The metric used for ranking impacts to the water column is the area of water surface in mi² that has been contaminated by 1 part per billion (ppb) oil to a depth of 33 feet. At 1 ppb, there are likely to be impacts to sensitive organisms in the water column and potential tainting of seafood, so this concentration is used as a screening threshold for both the ecological and socio-economic risk factors for water column resource impacts. To assist planners in understanding the scale of potential impacts for different leakage volumes, a regression curve was generated for the water column volume oiled using the five volume scenarios, which is shown in Figure 2-2, which is the regression curve for the *Cherokee*. Using this figure, the water column impacts can be estimated for any spill volume. On Figure 2-2, arrows are used to indicate the where the WCD for the *Taborfjell* plots on the curve and how the area of the water column impact is determined.

² French, D., M. Reed, K. Jayko, S. Feng, H. Rines, S. Pavignano, T. Isaji, S. Puckett, A. Keller, F. W. French III, D. Gifford, J. McCue, G. Brown, E. MacDonald, J. Quirk, S. Natzke, R. Bishop, M. Welsh, M. Phillips and B.S. Ingram, 1996. The CERCLA type A natural resource damage assessment model for coastal and marine environments (NRDAM/CME), Technical Documentation, Vol. I - V. Office of Environmental Policy and Compliance, U.S. Dept. Interior, Washington, DC.




Figure 2-2: Regression curve for estimating the area of water column at or above 1 ppb aromatics impacted as a function of spill volume for the *Taborfjell*. This regression curve was generated for the *Cherokee*, which has the same oil type and similar volume of potential releases as the *Taborfjell*. The arrows indicate where the WCD for the *Taborfjell* falls on the curve and how the area of water column impact can be determined for any spill volume.

Potential Water Surface Slick

The slick size from an oil release is a function of the quantity released. The estimated water surface coverage by a fresh slick (the total water surface area “swept” by oil over time) for the various scenarios is shown in Table 2-3, as the mean result of the 200 model runs for the *Cherokee* then using the regression curve shown in Figure 2-3 to calculate the values for the different release scenarios for the *Taborfjell*. Note that this is an estimate of total water surface affected over a 30-day period. In the model, the representative heavy fuel oil used for this analysis spreads to a minimum thickness of approximately 975 g/m^2 , and the oil is not able to spread any thinner, owing to its high viscosity. Thus, the results for the slick area swept are identical for the 0.01 and 10 g/m^2 thresholds. The slick will not be continuous but rather be broken and patchy. Surface expression is likely to be in the form of sheens, tarballs, and streamers. The location, size, shape, and spread of the oil slick(s) from an oil release from the *Taborfjell* will depend on environmental conditions, including winds and currents, at the time of release and in its aftermath. Refer to the risk assessment package for the *Cherokee* for maps (Figs. 2-2 and 2-3) showing the areas potentially affected by slicks using the Most Probable volume and the socio-economic and ecological thresholds.

Table 2-3: Estimated slick area swept on water for oil release scenarios from the *Taborfjell*, based on the model results for the *Cherokee*.

Scenario Type	Oil Volume (bbl)	Estimated Slick Area Swept Mean of All Models	
		0.01 g/m ²	10 g/m ²
Chronic	3	130 mi ²	130 mi ²
Episodic	30	410 mi ²	410 mi ²
Most Probable	300	1,300 mi ²	1,300 mi ²
Large	1,500	3,100 mi ²	3,100 mi ²
Worst Case Discharge	3,000	4,400 mi ²	4,400 mi ²

The actual area affected by a release will be determined by the volume of leakage, whether it is from one or more tanks at a time. To assist planners in understanding the scale of potential impacts for different leakage volumes, a regression curve was generated for the water surface area oiled using the five volume scenarios for the *Cherokee*, which is shown in Figure 2-3 and referenced in Table 2-3. Using this figure, the area of water surface with a barely visible sheen can be estimated for any spill volume from the *Taborfjell*.




Figure 2-3: Regression curve for estimating the amount of water surface oiling as a function of spill volume for the *Taborfjell*, showing both the ecological threshold of 10 g/m² and socio-economic threshold of 0.01 g/m², based on the model results for the *Cherokee*. The arrows indicate where the WCD for the *Taborfjell* falls on the curve and how the area of water surface impact can be determined for any spill volume. The curves for each threshold are so similar that they plot on top of each other.

Potential Shoreline Impacts

Based on these modeling results, shorelines from as far north as Maryland to as far south as Cape Canaveral, Florida are at risk. (Refer to Figure 2-6 in the *Cherokee* package to see the probability of oil stranding on the shoreline at concentrations that exceed the threshold of 1 g/m², for the Most Probable release). However, the specific areas that would be oiled will depend on the currents and winds at the time of the oil release(s), as well as on the amount of oil released. Estimated miles of shoreline oiling above the socio-economic threshold of 1 g/m² and the ecological threshold of 100 g/m² by scenario type are shown in Table 2-4.

Table 2-4: Estimated shoreline oiling from leakage from the *Taborfjell*, based on the modeling results for the *Cherokee*.

Scenario Type	Volume (bbl)	Estimated Miles of Shoreline Oiling Above 1 g/m ²	Estimated Miles of Shoreline Oiling Above 100 g/m ²
Chronic	3	6	0
Episodic	30	14	1
Most Probable	300	21	9
Large	1,500	27	14
Worst Case Discharge	3,000	29	17

The actual shore length affected by a release will be determined by the volume of leakage and environmental conditions during an actual release. To assist planners in scaling the potential impact for different leakage volumes, a regression curve was generated for the total shoreline length oiled using the five volume scenarios for the *Cherokee*, as detailed in Table 2-4 and shown in Figure 2-4. Using this figure, the shore length oiled can be estimated for any spill volume from the *Taborfjell*.




Figure 2-4: Regression curve for estimating the amount of shoreline oiling at different thresholds as a function of spill volume for the *Taborfjell*, based on the model results for the *Cherokee*. The arrows indicate where the WCD for the *Taborfjell* falls on the curve and how the length of shoreline impact can be determined for any spill volume.

SECTION 3: ECOLOGICAL RESOURCES AT RISK

Ecological resources at risk from a catastrophic release of oil from the *Taborfjell* (Table 3-1) include numerous guilds of birds that use shorelines and coastal waters. The islands of coastal Maine support an incredible diversity and abundance of nesting seabirds, migrating shorebirds and passerines and overwintering waterfowl. Shorelines in this region are important haul-out and pupping sites for seals. Coastal waters are summer foraging habitat for several species of large whales. Nearshore regions also support productive commercial fisheries for fish and invertebrate species.

Table 3-1: Ecological resources at risk from a release of oil from the *Taborfjell*.

(FT = Federal threatened; FE = Federal endangered; ST = State threatened; SE = State endangered).

Species Group	Species Subgroup and Geography	Seasonal Presence
Birds	<p><i>Important areas</i></p> <ul style="list-style-type: none"> • Digby Neck, NE Coastal Maine, Grand Manan Island, and Cape Cod region important for migratory songbirds and shorebirds • Coastal Maine is important breeding habitat for arctic tern (ST) • Over 365 of Maine's coastal islands have recent records of seabird nesting • Rocky islands of Maine, Nova Scotia, and New Brunswick are important wintering habitat for harlequin duck (ST) • Nearshore waters of Maine, Nova Scotia and New Brunswick are important habitat for razorbill (ST) and Atlantic puffin (ST) • Bay of Fundy mudflats are critical red knot stopover points • Shoreline from Casco Bay south is critical to least tern (FE, SE) and piping plover (FT, SE) nesting • Nantucket Sound, pelagic waters <4 km from land and offshore shoals and banks are concentration areas for sea birds and waterfowl • Cape Cod is a nationally significant migratory stopover site for numerous species, including red knot • Nearshore Cape Cod waters are important fall foraging areas for roseate tern (FE, SE) • Monomoy NWR and South Beach are the most important habitats in New England for nesting piping plover, American oystercatchers, and support major late-summer concentrations of shorebirds and roseate terns <p><i>Massachusetts nesting sites (numbers are in pairs unless otherwise noted)</i></p> <ul style="list-style-type: none"> • Cape Cod National Seashore: black skimmer, herring gull, great black-backed gull • Nantucket: herring gull, great black-backed gull, black-crowned night-heron, historic colonial waterbird nesting site, least tern, snowy egret, great egret • Nauset Marsh: roseate tern, common tern, black skimmer, arctic tern, laughing gull • Provincetown Harbor: double-crested cormorant, herring gull, great black-backed gull, common tern, black-crowned night-heron • Plymouth area: gulls, double-crested cormorant, least tern • Cohasset area: great black-backed gull, herring gull, double-crested cormorant • Boston Harbor Islands: American black duck, common eider, black-crowned night heron, glossy ibis, great egret, herring gull, great black-backed gull, double-crested cormorants Stratton Island: common tern (960), roseate tern, least tern (59), American oystercatcher (2) <p><i>Maine nesting sites</i></p>	<p>Nesting:</p> <p>Arctic tern nests May-Jun</p> <p>Razorbill present Mar-Sep; nests May-Jun</p> <p>Atlantic puffin nests Apr-Aug</p> <p>Least tern, roseate tern, common tern, piping plover present and nesting May-Aug</p> <p>Double-crested cormorant nests Apr-Jul</p> <p>Generally, seabird nesting season Apr-Aug</p> <p>Wintering:</p> <p>Harlequin ducks present Oct-Mar</p> <p>Migratory:</p> <p>Red knot present Jun-Jul</p> <p>Shorebird migration Aug</p> <p>Most summer residents leave by Sep</p> <p>Fall waterfowl migration Sep-Oct</p>

Species Group	Species Subgroup and Geography	Seasonal Presence
	<ul style="list-style-type: none"> Outer Green Island: common tern (1,067), arctic and roseate terns, black guillemots, common eiders Jenny Island: common tern (753), small numbers of roseate and arctic terns, common eiders Eastern Egg Rock: Atlantic puffin (123), roseate tern (90), common tern (829 pairs), laughing gull (2,051), Arctic tern, black guillemot, Leach's storm-petrel Stratton Island: northern limit of wading bird nesting (glossy ibis, little blue heron, great egret, tri-colored heron, American oystercatcher, least tern), southern limit of arctic bird nesting (common eider, black guillemot, arctic tern) 96% of Maine's population of roseate tern (FE) nest at Eastern Egg Rock and Stratton Island Maine Coastal Islands NWR: seabirds, waterfowl, wading birds, bald eagle, eiders, Arctic tern nesting, including razorbill (4 colonies) and Atlantic puffin (4 colonies): <ul style="list-style-type: none"> Petit Manan Island has 8 species of nesting seabirds, including Leach's storm-petrel and black guillemot Seal Island NWR: Atlantic puffin (546), arctic tern (1,201), common tern (1,836), razorbill, black guillemot, Leach's storm-petrel, occasionally roseate tern, common eider, gulls, and cormorants Matinicus rock: most diverse nesting colony on the U.S. Atlantic coast and only colony of manx shearwater; nesting species include Arctic tern (859), manx shearwater (4), puffins, razorbill, black guillemot, Leach's storm-petrel, common and roseate terns, laughing gull, common eider Pond Island NWR: common tern, occasionally roseate and arctic terns, common eider <p><i>Canada nesting sites</i></p> <ul style="list-style-type: none"> Sable Island: Leach's storm petrel, mallard, northern pintail, red-breasted merganser, black duck, spotted sandpiper, least sandpiper, herring and great black-backed gull, common and Arctic tern Grand Manan Archipelago: common eider, herring gull, black guillemot and great black-backed gull (common), common tern (1 colony), double-crested cormorant (3 colonies), razorbill and common murre (1 colony each), Leach's storm-petrel (5 colonies) 	
Pinnipeds	<p>Harbor seals and gray seals are common. Harp seals and hooded seals are transitory. Additional arctic species have been sighted but are not common.</p> <p><i>United States (Massachusetts and Maine):</i></p> <ul style="list-style-type: none"> High concentrations of harbor seals can be found at Monomoy NWR, Nauset marsh, Herring cove beach, Plymouth harbor, and the islands off Cohasset 30-40,000 harbor seals in coastal Maine and Isle of Shoals population Approx. 1,500-2,000 gray seals pup at Green Island, ME, Seal Island ME Gray seals pup at Cape Cod and Muskeget Island Harbor and Gray seals in high abundance in Jericho and Blue Hill Bay (ME) <p><i>Canada: Gray seal are more common than in U.S. waters.</i></p> <ul style="list-style-type: none"> Resident seal colonies on Grand Manan, Brier Island, and Sable Island Sable Island is an important pupping site for gray seals Seals common along SW shore of Nova Scotia Walrus can be found as far south as Sable Island 	<p>Harp seals present Jan-May</p> <p>Harbor seals pup May-Jun</p> <p>Gray seals pup Dec-Feb</p>
Cetaceans	<p><i>Common to Gulf of Maine/Bay of Fundy:</i> North Atlantic right whale (FE), fin whale (FE), minke whale, humpback whale (FE), Atlantic white-sided dolphin, harbor</p>	<p>Right, humpback, fin present in summer</p>

Section 3: Ecological Resources at Risk

Species Group	Species Subgroup and Geography	Seasonal Presence
	<p>porpoise</p> <ul style="list-style-type: none"> Cape Cod Bay is critical right whale foraging habitat Harbor porpoise more common in Bay of Fundy during the summer and move towards Cape Cod in the winter <p><i>Also present:</i> blue whale (FE), sei whale, pygmy sperm whale, sperm whale (FE), northern bottlenose whale, beaked whale, beluga, killer whale, long-finned pilot whale, white-beaked dolphin, bottlenose dolphin, common dolphin, striped dolphin</p> <p>Harbor porpoise more common in Bay of Fundy during the summer and move towards Cape Cod in the winter</p>	<p>Atlantic white-sided dolphin calve Jun-Jul</p> <p>Harbor porpoise calve May-Jun</p>
Fish	<p><i>Anadromous fish</i></p> <ul style="list-style-type: none"> Shortnose sturgeon (FE) found in Merrimack, Sheepscot, Kennebec, Androscoggin, and Penobscot Rivers, and Merrymeeting Bay Atlantic salmon (FE) spawn in streams north of Casco Bay, Maine Atlantic sturgeon (FT) spawn in Merrimack, Kennebec and Penobscot Rivers Alewife and American shad in all streams Coastal waters of Massachusetts are important areas for sturgeon, alewife Juvenile anadromous fish use nearshore waters as nursery habitat American shad concentrate in Northern Gulf of Maine in fall <p><i>Marine</i></p> <ul style="list-style-type: none"> Nearshore waters support high concentrations of anadromous fish Offshore ledges and banks support highly productive fisheries, including Atlantic halibut, Atlantic cod, winter flounder, witch flounder, American plaice, hake, monkfish Atlantic cod hotspots in nearshore waters of MA 	<p>Atlantic salmon spawn Jun-Apr</p> <p>Sturgeon spawn Apr-May</p> <p>American shad spawn May-Nov</p> <p>Alewife common offshore in the fall, spawn Mar-May</p>
Invertebrates	<p>Atlantic surf clam, softshell clam, bay scallop, northern quahog and blue mussels are all present in bays and nearshore environments. American lobster, rock crabs, northern shrimp and sea scallop beds all found in nearshore areas. Areas of high concentration are listed below:</p> <ul style="list-style-type: none"> Softshell clam, bay scallop, northern quahog in Pleasant Bay Blue mussels around Manomet Point Sea scallops offshore of Indian Brook conservation area Quahog and oyster in Cape Cod Bay Softshell clam beds common in bays of Maine American lobsters migrate inshore in the summer; rocky intertidal areas may be important nursery habitat <p>Northeastern beach tiger beetle (FT, SE) populations nest in the surf zone at Monomoy NWR and on Cape Cod</p>	<p>Northern shrimp Nov-May</p> <p>Mussels spawn in spring</p> <p>Beetles nest Jun-Sep</p>
Benthic Habitats	<p>Submerged aquatic vegetation (mostly eelgrass) is critical to numerous species and occurs inside of bays and sounds throughout the region</p> <p>Nantucket Harbor, Pleasant Bay, Plymouth Bay, and Casco Bay have large beds</p> <p>Common in protected environments throughout the region of impact</p> <p>Rockweed can be found along rocky shores from Maine north and is important habitat for juvenile fish, invertebrates and birds</p>	<p>Year round</p>

The Environmental Sensitivity Index (ESI) atlases for the potentially impacted coastal areas from a leak from the *Taborfjell* are generally available at each U.S. Coast Guard Sector. They can also be downloaded at: <http://response.restoration.noaa.gov/esi>. These maps show detailed spatial information on the

distribution of sensitive shoreline habitats, biological resources, and human-use resources. The tables on the back of the maps provide more detailed life-history information for each species and location. The ESI atlases should be consulted to assess the potential environmental resources at risk for specific spill scenarios. In addition, the Geographic Response Plans within the Area Contingency Plans prepared by the Area Committee for each U.S. Coast Guard Sector have detailed information on the nearshore and shoreline ecological resources at risk and should be consulted.

Ecological Risk Factors

Risk Factor 3: Impacts to Ecological Resources at Risk (EcoRAR)

Ecological resources include plants and animals (e.g., fish, birds, invertebrates, and mammals), as well as the habitats in which they live. All impact factors are based on a Worst Case and the Most Probable Discharge oil release from the wreck. Risk factors for ecological resources at risk (EcoRAR) are divided into three categories:

- Impacts to the water column and resources in the water column;
- Impacts to the water surface and resources on the water surface; and
- Impacts to the shoreline and resources on the shoreline.

The impacts from an oil release from the wreck would depend greatly on the direction in which the oil slick moves, which would, in turn, depend on wind direction and currents at the time of and after the oil release. Impacts are characterized in the risk analysis based on the likelihood of any measurable impact, as well as the degree of impact that would be expected if there is an impact. The measure of the degree of impact is based on the median case for which there is at least some impact. The median case is the “middle case” – half of the cases with significant impacts have less impact than this case, and half have more.

For each of the three ecological resources at risk categories, risk is defined as:

- The **probability of oiling** over a certain threshold (i.e., the likelihood that there will be an impact to ecological resources over a certain minimal amount); and
- The **degree of oiling** (the magnitude or amount of that impact).

As a reminder, the ecological impact thresholds are: 1 ppb aromatics for water column impacts; 10 g/m² for water surface impacts; and 100 g/m² for shoreline impacts.

In the following sections, the definition of low, medium, and high for each ecological risk factor is provided. Also, the classification for the *Taborfjell* is provided, both as text and as shading of the applicable degree of risk bullet, for the WCD release of 3,000 bbl and a border around the Most Probable Discharge of 300 bbl. Please note: The probability of oiling cannot be determined using the regression curves; probability can only be determined from the 200 model runs. Thus, the modeling results and regression curves for the *Cherokee* are used to estimate the values used in the risk scoring for the degree of oiling only.

Risk Factor 3A: Water Column Impacts to EcoRAR

Water column impacts occur beneath the water surface. The ecological resources at risk for water column impacts are fish, marine mammals, and invertebrates (e.g., shellfish, and small organisms that are food for larger organisms in the food chain). These organisms can be affected by toxic components in the oil. The threshold for water column impact to ecological resources at risk is a dissolved aromatic hydrocarbons concentration of 1 ppb (i.e., 1 part total dissolved aromatics per one billion parts water). Dissolved aromatic hydrocarbons are the most toxic part of the oil. At this concentration and above, one would expect impacts to organisms in the water column.

Risk Factor 3A-1: Water Column Probability of Oiling of EcoRAR (not scored)

This risk factor reflects the probability that at least 0.2 mi² of the upper 33 feet of the water column would be contaminated with a high enough concentration of oil to cause ecological impacts. The three risk scores for water column oiling probability are:

- **Low Oiling Probability:** Probability = <10%
- **Medium Oiling Probability:** Probability = 10 – 50%
- **High Oiling Probability:** Probability > 50%

Risk Factor 3A-2: Water Column Degree of Oiling of EcoRAR

The degree of oiling of the water column reflects the total volume of water that would be contaminated by oil at a concentration high enough to cause impacts. The three categories of impact are:

- | |
|---|
| <ul style="list-style-type: none"> • Low Impact: impact on less than 0.2 mi² of the upper 33 feet of the water column at the threshold level |
|---|
- **Medium Impact:** impact on 0.2 to 200 mi² of the upper 33 feet of the water column at the threshold level
 - **High Impact:** impact on more than 200 mi² of the upper 33 feet of the water column at the threshold level

The *Taborfjell* is classified as Low Risk for degree of oiling for water column ecological resources for the WCD of 3,000 bbl because the mean volume of water contaminated in the model runs was 0 mi² of the upper 33 feet of the water column. For the Most Probable Discharge of 300 bbl, the *Taborfjell* is also classified as Medium Risk for degree of oiling because the mean volume of water contaminated was 0 mi² of the upper 33 feet of the water column.

Risk Factor 3B: Water Surface Impacts to EcoRAR

Ecological resources at risk at the water surface include surface feeding and diving sea birds, sea turtles, and marine mammals. These organisms can be affected by the toxicity of the oil as well as from coating with oil. The threshold for water surface oiling impact to ecological resources at risk is 10 g/m² (10 grams of floating oil per square meter of water surface). At this concentration and above, one would expect impacts to birds and other animals that spend time on the water surface.

Risk Factor 3B-1: Water Surface Probability of Oiling of EcoRAR (not scored)

This risk factor reflects the probability that at least 1,000 mi² of the water surface would be affected by enough oil to cause impacts to ecological resources. The three risk scores for oiling are:

- **Low Oiling Probability:** Probability = <10%

- **Medium Oiling Probability:** Probability = 10 – 50%
- **High Oiling Probability:** Probability > 50%

Risk Factor 3B-2: Water Surface Degree of Oiling of EcoRAR

The degree of oiling of the water surface reflects the total amount of oil that would affect the water surface in the event of a discharge from the vessel. The three categories of impact are:

- **Low Impact:** less than 1,000 mi² of water surface impact at the threshold level
- **Medium Impact:** 1,000 to 10,000 mi² of water surface impact at the threshold level
- **High Impact:** more than 10,000 mi² of water surface impact at the threshold level

The *Taborfjell* is classified as Medium Risk for degree of oiling for water surface ecological resources for the WCD because the mean area of water contaminated in the model runs was 4,400 mi². It is also classified as Medium Risk for degree of oiling for the Most Probable Discharge because the mean area of water contaminated was 1,300 mi².

Risk Factor 3C: Shoreline Impacts to EcoRAR

The impacts to different types of shorelines vary based on their type and the organisms that live on them. For the modeled wrecks, shorelines were weighted by their degree of sensitivity to oiling. Wetlands are the most sensitive (weighted as “3” in the impact modeling), rocky and gravel shores are moderately sensitive (weighted as “2”), and sand beaches (weighted as “1”) are the least sensitive to ecological impacts of oil. In this risk analysis for the *Taborfjell*, shorelines have NOT been weighted by their degree of sensitivity to oiling because these data are available only for modeled vessels. Therefore, the impacts are evaluated only on the total number of shoreline miles oiled as determined from the regression curve.

Risk Factor 3C-1: Shoreline Probability of Oiling of EcoRAR (not scored)

This risk factor reflects the probability that the shoreline would be coated by enough oil to cause impacts to shoreline organisms. The threshold for shoreline oiling impacts to ecological resources at risk is 100 g/m² (i.e., 100 grams of oil per square meter of shoreline). The three risk scores for oiling are:

- **Low Oiling Probability:** Probability = <10%
- **Medium Oiling Probability:** Probability = 10 – 50%
- **High Oiling Probability:** Probability > 50%

Risk Factor 3C-2: Shoreline Degree of Oiling of EcoRAR

The degree of oiling of the shoreline reflects the length of shorelines oiled by at least 100 g/m² in the event of a discharge from the vessel. The three categories of impact are:

- **Low Impact:** less than 10 miles of shoreline impacted at the threshold level
- **Medium Impact:** 10 - 100 miles of shoreline impacted at the threshold level
- **High Impact:** more than 100 miles of shoreline impacted at the threshold level

The *Taborfjell* is classified as Medium Risk for degree of oiling for shoreline ecological resources for the WCD because the mean length of shoreline contaminated in the model runs was 17 miles. It is classified as Low Risk for degree of oiling for the Most Probable Discharge because the mean length of shoreline contaminated in the model runs was 9 miles.

Considering the modeled risk scores and the ecological resources at risk, the ecological risk from potential releases of the WCD of 3,000 bbl of heavy fuel oil from the *Taborfjell* is summarized as listed below and indicated in the far-right column in Table 3-2:

- Water column resources – Low, because little-to-no exposures above thresholds are likely
- Water surface resources – Medium, because of the seasonally very large number of marine mammal haulouts and nesting and migratory birds that use ocean, coastal, and estuarine habitats at risk. It should be noted that oil on the surface will not be continuous but rather be broken and patchy and in the form of sheens, tarballs, and streamers
- Shoreline resources – Medium, because of the many marine mammal haulouts and bird nesting areas associated with shorelines potentially at risk

Table 3-2: Ecological risk factor scores for the **Worst Case Discharge of 3,000 bbl** of heavy fuel oil from the *Taborfjell*.

Risk Factor	Risk Score			Explanation of Risk Score	Final Score
3A-1: Water Column Probability EcoRAR Oiling	Low	Medium	High	N/A: Only available for modeled vessels	Low
3A-2: Water Column Degree EcoRAR Oiling	Low	Medium	High	The mean volume of water contaminated above 1 ppb was 0 mi ² of the upper 33 feet of the water column	
3B-1: Water Surface Probability EcoRAR Oiling	Low	Medium	High	N/A: Only available for modeled vessels	Med
3B-2: Water Surface Degree EcoRAR Oiling	Low	Medium	High	The mean area of water contaminated above 10 g/m ² was 4,400 mi ²	
3C-1: Shoreline Probability EcoRAR Oiling	Low	Medium	High	N/A: Only available for modeled vessels	Med
3C-2: Shoreline Degree EcoRAR Oiling	Low	Medium	High	The length of shoreline contaminated by at least 100 g/m ² was 17 mi	

For the Most Probable Discharge of 300 bbl of heavy fuel oil, the ecological risk from potential releases from the *Taborfjell* is summarized as listed below and indicated in the far-right column in Table 3-3:

- Water column resources – Low, because little-to-no exposures above thresholds are likely
- Water surface resources – Medium, because the area affected is smaller, but there are still a large number of birds and marine mammals at risk. It should be noted that oil on the surface will not be continuous but rather be broken and patchy and in the form of sheens, tarballs, and streamers
- Shoreline resources – Medium, because of the many marine mammal haulouts and bird nesting areas associated with shorelines potentially at risk

Table 3-3: Ecological risk factor scores for the **Most Probable Discharge of 300 bbl** of heavy fuel oil from the *Taborfjell*.

Risk Factor	Risk Score			Explanation of Risk Score	Final Score
3A-1: Water Column Probability EcoRAR Oiling	Low	Medium	High	N/A: Only available for modeled vessels	Low
3A-2: Water Column Degree EcoRAR Oiling	Low	Medium	High	The mean volume of water contaminated above 1 ppb was 0 mi ² of the upper 33 feet of the water column	
3B-1: Water Surface Probability EcoRAR Oiling	Low	Medium	High	N/A: Only available for modeled vessels	Med
3B-2: Water Surface Degree EcoRAR Oiling	Low	Medium	High	The mean area of water contaminated above 10 g/m ² was 1,300 mi ²	
3C-1: Shoreline Probability EcoRAR Oiling	Low	Medium	High	N/A: Only available for modeled vessels	Med
3C-2: Shoreline Degree EcoRAR Oiling	Low	Medium	High	The length of shoreline contaminated by at least 100 g/m ² was 9 mi	

SECTION 4: SOCIO-ECONOMIC RESOURCES AT RISK

In addition to natural resource impacts, spills from sunken wrecks have the potential to cause significant social and economic impacts. Socio-economic resources potentially at risk from oiling are listed in Table 4-1 and shown in Figures 4-1 and 4-2. The potential economic impacts include disruption of coastal economic activities such as commercial and recreational fishing, boating, vacationing, commercial shipping, and other activities that may become claims following a spill.

Socio-economic resources in the areas potentially affected by a release from the *Taborfjell* include recreational beaches from Cape Cod and Nantucket, Massachusetts, up to northern Maine that are very highly utilized during summer, and are still in use during spring and fall for shore fishing. A national seashore and national park would also potentially be impacted.

A release could impact shipping lanes, which accommodate the ports of Boston, MA, Portsmouth, NH, Portland, ME, and Searsport, ME, totaling over 1,080 vessel calls and 57 million tonnage annually. Commercial fishing is economically important to the region. A release could impact fishing fleets coming out of several coastal towns and cities with annual catches totaling \$184.3 million.

In addition to the ESI atlases, the Geographic Response Plans within the Area Contingency Plans prepared by the Area Committee for each U.S. Coast Guard Sector have detailed information on important socio-economic resources at risk.

Spill response costs for a release of oil from the *Taborfjell* would be dependent on volume of oil released and specific areas impacted. The specific shoreline impacts and spread of the oil would determine the response required and the costs for that response.

Table 4-1: Socio-economic resources at risk from a release of oil from the *Taborfjell*.

Resource Type	Resource Name	Economic Activities
Tourist Beaches	Nantucket Island, MA Martha's Vineyard, MA Provincetown, MA Sagamore Beach, MA White Horse Beach, MA Priscilla Beach, MA Wellfleet, MA Truro, MA Duxbury, MA Chatham, MA Vinalhaven, ME Frenchboro, ME Cranberry Isles, ME Bar Harbor, ME Winter Harbor, ME Jonesport, ME Bar Harbor, ME Cutler, ME Truro, MA	Potentially affected beach resorts and beach-front communities in Massachusetts, New Hampshire, and Maine provide recreational activities (e.g., swimming, boating, recreational fishing, wildlife viewing, nature study, sports, dining, camping, and amusement parks) with substantial income for local communities and state tax income. Much of the coast is lined with economically-valuable beach resorts and residential communities. Many of these recreational activities are limited to or concentrated into the late spring through the early fall months.

Resource Type	Resource Name	Economic Activities
	Wellfleet, MA Provincetown, MA Chatham, MA	
National Seashores	Cape Cod National Seashore, MA	National seashores provide recreation for local and tourist populations while preserving and protecting the nation's natural shoreline treasures. National seashores are coastal areas federally designated as being of natural and recreational significance as a preserved area.
National Parks	Acadia National Park Boston Harbor Islands National Recreation Area	National parks also provide recreation for local and tourist populations and preserve and protect the nation's natural treasures.
National Wildlife Refuges	Nomans Land Island NWR (MA) Nantucket Island NWR (MA) Mashpee NWR (MA) Monomoy NWR (MA) Massasoit NWR (MA) Thacher Island NWR (MA) Great Bay NWR (NH) Rachel Carson NWR (ME) Pond Island NWR (ME) Franklin Island NWR (ME) Seal Island NWR (ME) Petit Manan NWR (ME) Cross Island NWR (ME)	National wildlife refuges in three states may be impacted. These federally managed and protected lands provide refuges and conservation areas for sensitive species and habitats.
State Parks	Birch Point SP, ME Camden Hills SP, ME Warren Island SP, ME Lamoine SP, ME Roque Bluffs SP, ME Quoddy Head SP, ME Hampton Beach SP, NH Jenness State Beach SP, NH Wallis Sands Beach SP, NH Ordiorne Point SP, NH	Coastal state parks are significant recreational resources for the public (e.g., swimming, boating, recreational fishing, wildlife viewing, nature study, sports, dining, camping, and amusement parks). They provide income to the states. State parks in the state of Maine are potentially impacted. Many of these recreational activities are limited to or concentrated into the late spring into early fall months.
Commercial Fishing	A number of fishing fleets use the area and surrounding waters for commercial fishing purposes.	
	Provincetown-Chatham, MA	Total Landings (2010): \$19.9M
	Gloucester, MA	Total Landings (2010): \$56.6M
	Portland, ME	Total Landings (2010): \$18.8M
	Rockland, ME	Total Landings (2010): \$10.6M
	Stonington, ME	Total Landings (2010): \$45.3M
	Boston, MA	Total Landings (2010): \$15.1M
	Jonesport, ME	Total Landings (2010): \$18.0M
Ports	There are a number of significant commercial ports in the Northeast that could potentially be impacted by spillage and spill response activities. The port call numbers below are for large vessels only. There are many more, smaller vessels (under 400 GRT) that also use these ports.	
	Boston, MA	584 port calls annually
	Portland, ME	317 port calls annually
	Searsport, ME	100 port calls annually
	Portsmouth, NH	83 port calls annually




Figure 4-1: Tribal lands, ports, and commercial fishing fleets at risk from a release from the *Taborfjell*. (Note that there are no tribal nations at risk.)




Figure 4-2: Beaches, coastal state parks, and Federal protected areas at risk from a release from the *Taborfjell*.

Socio-Economic Risk Factors

Risk Factor 4: Impacts to Socio-economic Resources at Risk (SRAR)

Socio-economic resources at risk (SRAR) include potentially impacted resources that have some economic value, including commercial and recreational fishing, tourist beaches, private property, etc. All impact factors are evaluated for both the Worst Case and the Most Probable Discharge oil release from the wreck. Risk factors for socio-economic resources at risk are divided into three categories:

- **Water Column:** Impacts to the water column and to socio-economic resources in the water column (i.e., fish and invertebrates that have economic value);
- **Water Surface:** Impacts to the water surface and resources on the water surface (i.e., boating and commercial fishing); and
- **Shoreline:** Impacts to the shoreline and resources on the shoreline (i.e., beaches, real property).

The impacts from an oil release from the wreck would depend greatly on the direction in which the oil slick moves, which would, in turn, depend on wind direction and currents at the time of and after the oil release. Impacts are characterized in the risk analysis based on the likelihood of any measurable impact, as well as the degree of impact that would be expected if there were one. The measure of the degree of impact is based on the median case for which there is at least some impact. The median case is the “middle case” – half of the cases with significant impacts have less impact than this case, and half have more.

For each of the three socio-economic resources at risk categories, risk is classified with regard to:

- The **probability of oiling** over a certain threshold (i.e., the likelihood that there will be exposure to socio-economic resources over a certain minimal amount known to cause impacts); and
- The **degree of oiling** (the magnitude or amount of that exposure over the threshold known to cause impacts).

As a reminder, the socio-economic impact thresholds are: 1 ppb aromatics for water column impacts; 0.01 g/m² for water surface impacts; and 1 g/m² for shoreline impacts.

In the following sections, the definition of low, medium, and high for each socio-economic risk factor is provided. Also, in the text classification for the *Taborfjell*, shading indicates the degree of risk for a WCD release of 3,000 bbl and a border indicates degree of risk for the Most Probable Discharge of 300 bbl. Please note: The probability of oiling cannot be determined using the regression curves; probability can only be determined from the 200 model runs. Thus, the modeling results and regression curves for the Cherokee are used to estimate the values used in the risk scoring for the degree of oiling only.

Risk Factor 4A-1: Water Column: Probability of Oiling of SRAR

This risk factor reflects the probability that at least 0.2 mi² of the upper 33 feet of the water column would be contaminated with a high enough concentration of oil to cause socio-economic impacts. The threshold for water column impact to socio-economic resources at risk is an oil concentration of 1 ppb (i.e., 1 part oil per one billion parts water). At this concentration and above, one would expect impacts

and potential tainting to socio-economic resources (e.g., fish and shellfish) in the water column; this concentration is used as a screening threshold for both the ecological and socio-economic risk factors.

The three risk scores for oiling are:

- **Low Oiling Probability:** Probability = <10%
- **Medium Oiling Probability:** Probability = 10 – 50%
- **High Oiling Probability:** Probability > 50%

Risk Factor 4A-2: Water Column Degree of Oiling of SRAR

The degree of oiling of the water column reflects the total amount of oil that would affect the water column in the event of a discharge from the vessel. The three categories of impact are:

- | |
|---|
| <ul style="list-style-type: none"> • Low Impact: impact on less than 0.2 mi² of the upper 33 feet of the water column at the threshold level |
|---|
- **Medium Impact:** impact on 0.2 to 200 mi² of the upper 33 feet of the water column at the threshold level
 - **High Impact:** impact on more than 200 mi² of the upper 33 feet of the water column at the threshold level

The *Taborfjell* is classified as Low Risk for degree of oiling for water column socio-economic resources for the WCD of 3,000 bbl because the mean volume of water contaminated in the model runs was 0 mi² of the upper 33 feet of the water column. For the Most Probable Discharge of 300 bbl, the *Taborfjell* is classified as Low Risk for degree of oiling because the mean volume of water contaminated 0 mi² of the upper 33 feet of the water column.

Risk Factor 4B-1: Water Surface Probability of Oiling of SRAR

This risk factor reflects the probability that at least 1,000 mi² of the water surface would be affected by enough oil to cause impacts to socio-economic resources. The three risk scores for oiling are:

- **Low Oiling Probability:** Probability = <10%
- **Medium Oiling Probability:** Probability = 10 – 50%
- **High Oiling Probability:** Probability > 50%

The threshold level for water surface impacts to socio-economic resources at risk is 0.01 g/m² (i.e., 0.01 grams of floating oil per square meter of water surface). At this concentration and above, one would expect impacts to socio-economic resources on the water surface.

Risk Factor 4B-2: Water Surface Degree of Oiling of SRAR

The degree of oiling of the water surface reflects the total amount of oil that would affect the water surface in the event of a discharge from the vessel. The three categories of impact are:

- **Low Impact:** less than 1,000 mi² of water surface impact at the threshold level
- | |
|---|
| <ul style="list-style-type: none"> • Medium Impact: 1,000 to 10,000 mi² of water surface impact at the threshold level |
|---|
- **High Impact:** more than 10,000 mi² of water surface impact at the threshold level

The *Taborfjell* is classified as Medium Risk for degree of oiling for water surface socio-economic resources for the WCD of 3,000 bbl because the mean area of water contaminated in the model runs was 4,400 mi². The *Taborfjell* is classified as Medium Risk for degree of oiling for water surface socio-economic resources for the Most Probable Discharge because the mean area of water contaminated was 1,300 mi².

Risk Factor 4C: Shoreline Impacts to SRAR

The impacts to different types of shorelines vary based on economic value. For the modeled wrecks, shorelines have been weighted by their degree of sensitivity to oiling. Sand beaches are the most economically valued shorelines (weighted as “3” in the impact analysis), rocky and gravel shores are moderately valued (weighted as “2”), and wetlands are the least economically valued shorelines (weighted as “1”). In this risk analysis for the *Taborfjell*, shorelines have NOT been weighted by their degree of sensitivity to oiling because these data are available only for modeled vessels. Therefore, the impacts are evaluated only on the total number of shoreline miles oiled as determined from the regression curve.

Risk Factor 4C-1: Shoreline Probability of Oiling of SRAR

This risk factor reflects the probability that the shoreline would be coated by enough oil to cause impacts to shoreline users. The threshold for impacts to shoreline SRAR is 1 g/m² (i.e., 1 gram of oil per square meter of shoreline). The three risk scores for oiling are:

- **Low Oiling Probability:** Probability = <10%
- **Medium Oiling Probability:** Probability = 10 – 50%
- **High Oiling Probability:** Probability > 50%

Risk Factor 4C-2: Shoreline Degree of Oiling of SRAR

The degree of oiling of the shoreline reflects the total amount of oil that would affect the shoreline in the event of a discharge from the vessel. The three categories of impact are:

- **Low Impact:** less than 10 miles of shoreline impacted at threshold level
- **Medium Impact:** 10 - 100 miles of shoreline impacted at threshold level
- **High Impact:** more than 100 miles of shoreline impacted at threshold level

The *Taborfjell* is classified as Medium Risk for degree of oiling because the mean length of shoreline contaminated in the model runs was 29 miles. The *Taborfjell* is classified as Medium Risk for degree of oiling for shoreline socio-economic resources for the Most Probable Discharge because the mean length of shoreline contaminated was 21 miles.

Considering the modeled risk scores and the socio-economic resources at risk, the socio-economic risk from potential releases of the WCD of 3,000 bbl of heavy fuel oil from the *Taborfjell* is summarized as listed below and indicated in the far-right column in Table 4-2:

- Water column resources – Low, because there would be virtually no impact on the water column
- Water surface resources – Low, because there would be a relatively small surface area offshore impacted in important shipping lanes. It should be noted that oil on the surface will not be continuous but rather be broken and patchy and in the form of sheens, tarballs, and streamers
- Shoreline resources – Medium, because a relatively small length of shoreline would be impact, although there are high-value shoreline resources at risk

Table 4-2: Socio-economic risk factor ranks for the **Worst Case Discharge of 3,000 bbl** of heavy fuel oil from the *Taborfjell*.

Risk Factor	Risk Score			Explanation of Risk Score	Final Score
4A-1: Water Column Probability SRAR Oiling	Low	Medium	High	N/A: Only available for modeled vessels	Low
4A-2: Water Column Degree SRAR Oiling	Low	Medium	High	The mean volume of water contaminated above 1 ppb was 0 mi ² of the upper 33 feet of the water column	
4B-1: Water Surface Probability SRAR Oiling	Low	Medium	High	N/A: Only available for modeled vessels	Low
4B-2: Water Surface Degree SRAR Oiling	Low	Medium	High	The mean area of water contaminated above 0.01 g/m ² was 4,400 mi ²	
4C-1: Shoreline Probability SRAR Oiling	Low	Medium	High	N/A: Only available for modeled vessels	Med
4C-2: Shoreline Degree SRAR Oiling	Low	Medium	High	The length of shoreline contaminated by at least 1 g/m ² was 29 mi	

For the Most Probable Discharge of 300 bbl, the socio-economic risk from potential releases of heavy fuel oil from the *Taborfjell* is summarized as listed below and indicated in the far-right column in Table 4-3:

- Water column resources – Low, because there would be virtually no impact on the water column
- Water surface resources – Low, because there would be a relatively small surface area offshore impacted in important shipping lanes. It should be noted that oil on the surface will not be continuous but rather be broken and patchy and in the form of sheens, tarballs, and streamers
- Shoreline resources – Medium, because a relatively small length of shoreline would be impact, although there are high-value shoreline resources at risk

Table 4-3: Socio-economic risk factor ranks for the **Most Probable Discharge of 300 bbl** of heavy fuel oil from the *Taborfjell*.

Risk Factor	Risk Score			Explanation of Risk Score	Final Score
4A-1: Water Column Probability SRAR Oiling	Low	Medium	High	N/A: Only available for modeled vessels	Low
4A-2: Water Column Degree SRAR Oiling	Low	Medium	High	The mean volume of water contaminated above 1 ppb was 0 mi ² of the upper 33 feet of the water column	
4B-1: Water Surface Probability SRAR Oiling	Low	Medium	High	N/A: Only available for modeled vessels	Low
4B-2: Water Surface Degree SRAR Oiling	Low	Medium	High	The mean area of water contaminated above 0.01 g/m ² was 1,300 mi ²	
4C-1: Shoreline Probability SRAR Oiling	Low	Medium	High	N/A: Only available for modeled vessels	Med
4C-2: Shoreline Degree SRAR Oiling	Low	Medium	High	The length of shoreline contaminated by at least 1 g/m ² was 21 mi	

SECTION 5: OVERALL RISK ASSESSMENT AND RECOMMENDATIONS FOR ASSESSMENT, MONITORING, OR REMEDIATION

The overall risk assessment for the *Taborfjell* is comprised of a compilation of several components that reflect the best available knowledge about this particular site. Those components are reflected in the previous sections of this document and are:

- Vessel casualty information and how site formation processes have worked on this vessel
- Ecological resources at risk
- Socio-economic resources at risk
- Other complicating factors (war graves, other hazardous cargo, etc.)

Table 5-1 summarizes the screening-level risk assessment scores for the different risk factors, as discussed in the previous sections. As noted in Sections 3 and 4, each of the ecological and socio-economic risk factors each has two components, probability and degree. Please note: The probability of oiling cannot be determined using the regression curves; probability can only be determined from the 200 model runs. Thus, the modeling results and regression curves for the *Cherokee* were used to estimate the values used in the risk scoring for the **degree of oiling only**.

In order to make the scoring more uniform and replicable between wrecks, a value was assigned to each of the 7 criteria. This assessment has a total of 7 criteria (based on table 5-1) with 3 possible scores for each criteria (L, M, H). Each was assigned a point value of L=1, M=2, H=3. The total possible score is 21 points, and the minimum score is 7. The resulting category summaries are:

Low Priority	7-11
Medium Priority	12-14
High Priority	15-21

For the Worst Case Discharge, *Taborfjell* scores Low with 11 points; for the Most Probable Discharge, *Taborfjell* also scores Low with 11 points. Under the National Contingency Plan, the U.S. Coast Guard and the Regional Response Team have the primary authority and responsibility to plan, prepare for, and respond to oil spills in U.S. waters. Based on the technical review of available information, NOAA proposes the following recommendations for the *Taborfjell*. The final determination of what type of action, if any, rests with the U.S. Coast Guard.

<i>Taborfjell</i>	Possible NOAA Recommendations
	Wreck should be considered for further assessment to determine the vessel condition, amount of oil onboard, and feasibility of oil removal action
	Use surveys of opportunity to attempt to locate this vessel and gather information on vessel condition
	Conduct active monitoring to look for releases or changes in rates of releases
✓	Be noted in the Area Contingency Plans so that if a mystery spill is reported in the general area, this vessel could be investigated as a source
✓	Conduct outreach efforts with the technical and recreational dive community as well as commercial and recreational fishermen who frequent the area, to gain awareness of changes in the site

This vessel is of historic significance and will require appropriate actions be taken under the National Historic Preservation Act and possibly the Sunken Military Craft Act prior to any actions that could

impact the integrity of the vessel. The site is also considered a war grave and appropriate actions should be undertaken to minimize disturbance to the site.

Table 5-1: Summary of risk factors for the *Taborfjell*.

Vessel Risk Factors		Data Quality Score	Comments	Risk Score	
Pollution Potential Factors	A1: Oil Volume (total bbl)	Low	Maximum of 3,000 bbl, not reported to be leaking	Med	
	A2: Oil Type	High	Bunker oil is heavy fuel oil, a Group IV oil type		
	B: Wreck Clearance	High	Vessel not reported as cleared		
	C1: Burning of the Ship	High	No fire was reported		
	C2: Oil on Water	High	No oil was reported on the water		
	D1: Nature of Casualty	Medium	One (possibly two torpedo detonations)		
	D2: Structural Breakup	Low	Unknown structural breakup		
Archaeological Assessment	Archaeological Assessment	Low	Limited sinking records of this ship exist, and the site has never been located, assessment is believed to have low accuracy	Not Scored	
Operational Factors	Wreck Orientation	Low	Unknown, potential to be upright	Not Scored	
	Depth	Low	> 800 ft		
	Visual or Remote Sensing Confirmation of Site Condition	Low	Location unknown		
	Other Hazardous Materials Onboard	High	No		
	Munitions Onboard	High	Munitions for onboard weapons		
	Gravesite (Civilian/Military)	High	Yes		
	Historical Protection Eligibility (NHPA/SMCA)	High	NHPA and possibly SMCA		
				WCD	Most Probable
Ecological Resources	3A: Water Column Resources	High	Little-to-no exposures above thresholds likely	Low	Low
	3B: Water Surface Resources	High	Seasonally very high concentrations of marine birds in coastal and shelf waters	Med	Med
	3C: Shore Resources	High	Sand beaches and rocky shores at risk include many marine mammal haulouts and bird nesting sites	Med	Med
Socio-Economic Resources	4A: Water Column Resources	High	Virtually no impact on the water column	Low	Low
	4B: Water Surface Resources	High	A relatively small to moderate surface area offshore impacted in important shipping lanes	Low	Low
	4C: Shore Resources	High	A relatively small length of shoreline would be impact, although there are high-value shoreline resources at risk	Med	Med
Summary Risk Scores				11	11